

The inherent potential of Algae for Forthcoming Future: a Comprehensive Review

Jemilakshmi.T.V^{1*}

^{1*}Assistant Professor, Faculty of Engineering, Department of Biotechnology, Karpagam Academy of Higher Education, Coimbatore, Tamil Nadu, India,

Corresponding Author: e-mail - jemilakshmi.tv@kahedu.edu.in

ABSTRACT

Algae are an extensive and varied community of microbes that can take out photosynthesis as they absorb sunlight. Blue-green algae are a well-known source of renewable energy and exhibit an enormous diversity in terms of morphology and metabolism. Algae, mainly marine algae theoretically have the potential as fertilizers. This could result in a lesser amount of nitrogen and phosphorous overflow when compared to the utilization of livestock fertilizers. This, successively, raises the quality of water. These organisms are grown all over the world and used as supplements for human nutriment. Algae have a vast perspective as a bioresource for different sectors like biofuels, pharmaceuticals, nutraceuticals, and the food industry. This interesting cluster of organisms also has relevance in cultivation. This can facilitate improved nutrient accessibility, keep up the organic carbon and fertility of the soil, and improving plant development and crop yields. Biofuels have emerged as a safe, nature-friendly, cost-effective alternative to other fuels. Many government agencies and companies are putting efforts for bringing down the capital and operating cost and thus make the production of algal biofuel commercially feasible. This review addresses applications of algae in agriculture, food, and other industries. It is found that further research work is required to optimize algae production and to assess the possibility of more environmentally friendly goods with higher added value.

KEYWORDS: Blue-green algae, Agriculture, Biofuel, Wastewater treatment, Food industry, Medicinal application.

1. INTRODUCTION

Algae are omnipresent and they grow in pretty much every livable environment on earth, in soils, ice, snowfields, underground aquifers, and warm and cold deserts. Biochemically and physiologically, algae are comparable in numerous viewpoints to other flora (Lee et al. 1989). The algae are of various shapes and sizes. The tiny algae are referred to as phytoplankton and perceptible ones as seaweeds (Dhargalkar and Pereira 2005). Because of the strange increment in algal biomass, especially that of green kelp, coming about because of the dynamic eutrophication of some coastal ecosystems, the interest in the agricultural use of seaweeds as composts have been increased in the last decades (Morand and Briand 1996). Recent studies have demonstrated that algae have an important role in securing plants against various biotic and abiotic stresses and offer the potential for field application. At present, the growing interest in natural strategies for crop production, as well as

the diminishing number of chemical products allowed for use in the fields, is rousing the quest for new bioformulations (Colavita et al. 2011).

In general, microalgae are categorized into Chlorophyta, Rhodophyta, Phaeophyta, Euglenophyta, Pyrrophyta, and Chrysophyta. The only photosynthetic prokaryotes able to generate oxygen is Cyanobacteria (Singh et al. 2015). There are around 30,000 species of both unicellular microalgae as well as progressively complex multicellular organisms (Rath et al. 2012).

The structural attributes of eukaryotic microalgae and prokaryotic cyanobacteria take after a thallus, i.e. not separated into roots or shoots. The filamentous cyanobacteria are divided into heterocystous and non-heterocystous structures. Most algae play a significant job in sequestering natural carbon and are liable for half of the whole photosynthesis process on the globe (Moroney and Ynalvez 2009). Besides, current studies have uncovered the possible utilization of green microalgae as organic fertilizer, for upgraded fertility of the soil, plant development, quality of the organic product, and nourishing attributes, and grain yield (Coppens et al. 2016). Microalgae, especially cyanobacteria are likewise measured as possible biocontrol operators as they show adversarial impact against several plant pathogens, for example, microscopic organisms, parasites, and nematodes, chiefly as an aftereffect of the creation of hydrolytic chemicals and biocidal mixes, for example, benzoic acid, majusculonic acid, etc (Chandel 2009; Chaudhary et al. 2012; Prasanna et al. 2008b; Gupta et al. 2013). Microalgae are alluring as fuel sources because numerous species grow relatively fast compared to terrestrial plants and can be developed on brackish or saline water, in this way avoiding the utilization of unsustainable amounts of freshwater (Dismukes et al. 2008). Microalgae also have potential applications in the field of the food industry, medical field, wastewater treatment, etc (Jayachitra et al, 2021).

2. APPLICATION OF ALGAE

2.1 In Agriculture:

Presently utilized conventional farming management practices intensely depend on the use of chemical composts and pesticides, and practices like rigorous tillage and excess irrigation which or else lead to expanding farming cost, excess abuse of soil and water, make pollution (Kumar et al. 2012).

Algae are proven for positive effects on the quality of the soil. Algae improve the fertility of the soil by expanding the general soil activity and improve microbial interactions as they maintain the development of useful microorganisms (Leloup et al. 2013; Thomas 2012; Yan-Gui et al. 2013).

Cyanobacteria are suitable organic fertilizers for rice-based cropping methods. It's a significant part of wetland rice ecosystems and serves as the least expensive source of natural biofertilizers (Omar 2000; Ladha and Reddy 2003). For carbon and nitrogen fixation diazotrophic cyanobacteria need daylight as a solitary energy resource. They have the incredible possibility of using as biofertilizers, and their utilization will diminish fuel requirement for fertilizer production. The agronomic capability of heterocystous cyanobacteria, both free-living or in a symbiotic relationship with *Azolla*, has been accepted (El-Zeky et al. 2005).

Inoculation of cyanobacteria in soil has been analyzed to support the plants in multiple points of view, for example, improving germination of seeds, growth of plants, grain yield, and nutritional

enhancement(Karthikeyan et al.2007; Prasanna et al.2016a; Prasanna et al.2016b).Cyanobacteria have some phosphate solubilizing variety. Phosphorus (P) is the second significant supplement after nitrogen for plant life and microbes. P and N are limiting supplements for most aquatic systems (Silke et al.2007).

While dealing with the algae of Indian rice fields, Gupta et al(2013) noticed that cyanobacteria quickened the germination of seedsand enhanced sprout development. Likewise, they additionally noticed that both the yield and the quality of the grains were enhanced in terms of protein content. The positive effect of the algae on the rice yield may not be limited to their capability to fix atmospheric nitrogen alone; also they have addedadvantageous roles like releasing bioactive matter.It is also clear that the co-inoculation of cyanobacteria with wheat improved root dry mass and chlorophyll content (Obreht et al.1993). In 1995 Gantaret al (1995)noticed that extracellular substances discharged by cyanobacteria that inhabiton roots of wheat plants indicated a noteworthy impact on plant growth. The quick cyanobacterial cell development and simple nutritional necessities primarily water, sunlight,and CO₂give a wide degree to the commercial utilization of cyanobacterial species as plant growth promoters(Ruffing 2011).

Algae are well-knownfor the presence of plant hormones (Tarakhovshaya et al.2007; Khan et al.2009).On the other hand, their concentration is not oftenanalyzed particularly in algal concentrates acquired by supercritical fluid extraction. The supercritical concentrate acquired from Baltic seaweeds consists of plant hormones, for example, phenylacetic corrosive (PAA) from the auxin group and 6-benzyl amino purine (6-BA) from the cytokinin group (Michalak et al.2016). Intheworkof Amin et al. [2009],several phytohormones were found in an aqueous extract of spirulina.

2.2 Biofuel Production

Biofuel refers to any kind of solid, liquid, or gaseous fuels.This can be resultantof renewable resources.Microalgal biomass for the biodiesel makingprocess has been done using either pure or mixed cultures. Autotrophic cultivation can be done in either open or closed systems and heterotrophic growth is kept in fermentors (Singh et al.2011).

Developing microalgae on various sorts of wastewaters like rural run-off, intenseanimal feedstocks, and waste streams have been analyzed over the previous years. The achievement of those investigations intensely relies upon the strains of microalgae. Numerous microalgae strains, for example, Chlorella sp., Scenedesmus sp., Micractinium sp., Actinastrum sp., Heynigia sp., Hindakia sp., Pediastrum sp.,etc have been tried and found to use for evacuating N and P components in the wastewaters (Woertz et al.2009; Wang et al.2010; Oswald et al.1957).For the production of biofuels, collectedlow-priced algal biomass could be used as the best input. There is a vast requirement for more healthystrains of microalgaethat are resistant to particular types of sewage water. Many researchers found that microalgae modified to various culture conditions can grow well than those obtained from algae banks Zhou et al.2012; Li et al.2010).

The steps involved in the biomass processing system are microalgae culturing, collection, dewatering, and drying. For the algal oil conversion process, two methods are generally following. A two-stage technique that is split intothe extraction of oil and oil transesterification and one step, in situ transesterification. Three kinds of conversiontechniques exist, chemical, thermochemical, and enzymatic (Maurycy et al.2013).

2.3 Wastewater Treatment

Microalgae are universally perceived for their capacity to expel minerals and heavy metals and thus, collect lipid and starch molecules. The fixation of Carbon, Nitrogen, and Phosphorous (C, N, and P) in sewage water empowers the development of microalgae on a higher amounts since they need 358, 63, and 9 mg L⁻¹ for delivering 1kg of algal biomass (Kong et al.2010). Algae can cultivate in sewage water by using nitrate and phosphate as supplements (Shobana et al.2017).

Cyanobacteria and microalgae are additionally useful in the deprivation of oil and oil mixes (Abed 2010; Abed and Koster 2005). Abed and Koster (2005) assessed five cyanobacteria- *Aphanothece halophyletica*, *Dactylococcopsis salina*, *Halothece* sp., *Oscillatoria* sp., and *Synechocystis* sp. fit for degradation of n-alkane. This study indicated that the deprivation was associated with petroleum degrading aerobic microbes (Abed 2010; Abed and Koster 2005).

Green microalga *Scenedesmus* sp. gave maximal algal biomass with 106 cells mL⁻¹ with half market sewage water (98.54 mg L⁻¹ day⁻¹) and high expulsion of total nitrogen (TN), phosphorous (TP), and total organic carbon (TOC) at 85, 90, and 65 % correspondingly (Apandi et al.2018).

Treatment of wastewater using microalgae not just decreases the contamination rate in the sewage but also considerably builds the biomass grouping of microalgae. This is significant to the research did by Ammar et al. (2018) where effluent treatment and biomass development by *Nannochloropsis oculata* and *Isochrysis galbana* in oilfield produced water (PW) were analyzed.

2.4 In Food Industry

Algae have been used as food for many years in diverse cultures (Wells et al., 2017). Archeological records in Chile show utilization of algae dating to 14,000 years back, This is marked as the most seasoned known utilization of algae (Dillehay et al., 2008). They can essentially fulfill the requirement for protein, particularly from the viewpoint of a rising populace. Microalgae protein (MP) can sometimes be superior to the present overwhelming protein sources, particularly in light of the fact that they need low land requirement for cultivation.

Because of the gelling and thickening properties, the foodstuffs like agar, alginates, and carrageenans are considered the most significant items which can be obtained from algae. In past years, noteworthy development has been seen in areas of algae research. Progress in various areas such as protoplast fusion, macroalgal cell cultures, transgenic algae are also notable (Bhattacharjee 2016). Alginate, a brown alga derivative is utilized in the textile business for sizing cotton yarn. It also has significance in the food industry because of its gelling nature. Alginate is well known for its chelating property and capacity to make an extremely thick solution. This makes it a good nominee for food and drug industries (Ververis et al.2007).

Seaweeds are utilized in human and animal nutrition. Customers are captivated towards particular variety, for example, *Porphyra* sp., *Chondrus crispus*, *Himanthalia elongate*, and *Undaria pinnatifida*, and towards the food industry as a result of their low calory content and high nutrients content, minerals and dietary fiber (Plaza et al.2008). Algal biomass is available as a powder, tablets, fluids, etc. it can be fused into various food items and the most significant algal varieties in human nutrition are *Spirulina* and *Chlorella* sp. Consumable algae are the main source of dietary fiber, minerals, and proteins (Kuda et al.2002).

Group/species	Extract	Use/application
<i>Aphanizomenon flos-aquae</i>	Protein, essential fatty acids, β -carotene	Health foods, food supplements
<i>Arthrospira maxima</i>	Omega 3 PUFAs	Health foods
<i>Arthrospira platensis</i>	Protein, fiber, mineral content.	Health foods
<i>Chlorella</i> spp.	Biomass, carbohydrate extract.	Animal nutrition, health drinks, food supplement
<i>Cryptocodinium cohnii</i>	DHA	Infant health and nutrition
<i>Diacronema vikianum</i>	Omega 3 PUFAs	Health foods
<i>Dunaliella</i> sp.	Protein content	Health foods
<i>Haematococcus pluvialis</i>	Carotenoids, astaxanthin	Health food, food supplement, feeds
<i>Himantalia elongate</i>	PUFAs, α -Tocopherol, Sterols, Soluble fibres.	Health food, food supplement
<i>Isochrysis galbana</i>	Omega 3 PUFAs	Health foods
<i>Schizochytrium</i> sp.	DHA and EPA	Food, beverage, and food supplement
<i>Ulva</i> spp	Sterols	Health foods

Table 1: Microalgae species suitable for food products (Arkadiusz and Joanna, 2020)

2.5 Medicinal Applications

One of the microalgae focused compound is a bioactive compound with anticancer activities. About 1000 extracts of various cyanobacteria sp. were screened out for antineoplastic property. *Potriochromonas malhamensis* has demonstrated inhibition of protein tyrosine kinase activity because of the existence of a novel chlorosulfolipid compound. Many cyanobacteria are promising producers of bioactive compounds. These can destroy cancer cells by the apoptosis process, affecting cell signaling by inducing protein kinase C family signaling enzymes. Preliminary studies have shown that certain antioxidants like β -Carotene, may be beneficial in the treatment of precancerous conditions (Boopathy and Kathiresan 2010).

Specific extracts from 600 Cyanobacteria cultures were analyzed for cellular inhibition of infection such as HIV-1 and HSV-2. But this shows only a 10 percent rate of success. Patterson et al., (1993) isolated a new cyanobacterial compound called cyanovirin-N, which is an effective virucidal agent against HIV. Cyanovirin-N can block the contact of viral glycoprotein gp120 with CD4 (Dev et al. 2000). Rinehart et al. (1981) stated the antiviral activity of 5% Cyanobacteria extracts against Herpes simplex virus type II. Under 5% of the extract has proven for activity in opposition to Respiratory syncytial virus (Patterson et al. 1991).

The microalgal cell has antibacterial activity both against gram-positive and gram-negative bacteria. There are also antifungal occurrences in green algae, diatoms, and dinoflagellates.

Microalgal toxins obtained from blue-green algae, *Ochromonas* sp., and *Prymnesium parvum* are applied in pharmaceutical sector (Katircioglu et al. 2006; Borowitzka et al. 1988). Various strains of Cyanobacteria generate intra and extracellular metabolites. These strains were also proven for antifungal, antialgal, antiviral, and antibacterial properties. Antimicrobial agent creation is influenced by several factors like incubation temperature, pH of culture medium, incubation period, light intensity, and medium composition (Noaman et al. 2004).

3. CONCLUSION

The widely used metaphoric concept of algal biomass is an “untapped resource” that suggests that natural products and related benefits to the economy and human well-being will flow until the correct valve is found and opened. Genetic and metabolic modifications of Cyanobacteria play significant roles in enhancing the economy of Cyanobacteria-mediated bio-fuel production in the future. This review highlighted the use of a variety of algae in limitless sectors or applications such as bioenergy, wastewater treatment, medical sector, food industry, and agriculture.

The effect of blue-green algae has been studied in paddy fields under flooded surroundings, which are perfect for the development of blue-green algae, but their impact on dry crops such as wheat, maize, etc. needs to be explored and will additionally improve their use in agriculture. Since algal biomass is a significant tool for developing environmentally responsible approaches to cultivation in addition to constituting the basis of food webs. There is a high need to raise attentiveness and encourage in-depth research work to expand their applications in the coming years.

The latest advances in the use of consortia or biofilms of green algae and cyanobacteria as biofertilizers have been proved to be promising. The use of algae for the development of biofuels provides many advantages compared with higher plants, i.e., least land for farming, fast growth rate, high productivity of lipids, and biomass. Furthermore, the full use of algae would create zero waste. Algae is an all-rounder in the medical, farming, and bioremediation sectors. The rich algal variety needs to be harnessed for various futile applications. In summary, algae are gifted sources of biofuels, high-value molecules, nutraceuticals, and a variety of bioactive metabolites for exploring out novel drugs in terms of their market applicability.

REFERENCE

1. Abed R.M.M. (2010) Interaction between cyanobacteria and aerobic heterotrophic bacteria in the degradation of hydrocarbons. *Int. Biodeterior. Biodegrad.* 64 (1), 58–64.
2. Abed R.M.M., & Koster J. (2005) The direct role of aerobic heterotrophic bacteria associated with cyanobacteria in the degradation of oil compounds. *Int. Biodeterior. Biodegrad.* 55 (1), 29–37.
3. Amin G.H., Al-Gendy A.A., Yassin El-Ayouty M., & Abdel-Motteleb A. (2009) Effect of *Spirulina platensis* extract on growth, phenolic compounds and antioxidant activities of *Sisymbrium Iriocallus* and cell suspension cultures. *Austr. J. Basic Appl. Sci.* 3: 2097–2110.

4. AmmarS.H., Khadim H.J.,& Mohamed A.I. (2018) Cultivation of *Nannochloropsis oculata* and *Isochrysis galbana* microalgae in produced water for bioremediation and biomass production. *Environ Technol Inno.* 10:132-142.
5. Apandi N., Mohamed R.M.S.R., Al-Gheethi A., Gani P., Ibrahim A., Kassim A.H.M. (2018) *Scenedesmus* Biomass Productivity and Nutrient Removal from Wet Market Wastewater. A Bio-kinetic Study. *c. 10(10): 2783–2800.*
6. Arkadiusz Piwowar and Joanna Harasym.(2020) The Importance and Prospects of the Use of Algae in Agribusiness. *Sustainability*, 12, 5669.
7. Bhattacharjee M. (2016) Pharmaceutically valuable bioactive compounds of algae. *Asian J Pharm Clin Res.* 9: 43-47.
8. BoopathyN.S.,&KathiresanK. (2010) Anticancer drugs from marine flora: An overview. *J.Oncol.* 4:1-18.
9. Borowitzka M. & Borowitzka L. (1988) *Micro-algal biotechnology.* New York, USA:Cambridge University press.
10. Chandel S.T. (2009) Nematicidal activity of the Cyanobacterium, *Aulosira fertilissima* on the hatch of *Meloidogynetritycoryzae* and *Meloidogyneincognita*. *Arch. Phytopathol. Plant Protect.* 42 (1): 32–38.
11. ChaudharyV., Prasanna R., Nain L., Dubey S.C., Gupta V., SinghR.,Jaggi S.,& BhatnagarA.K. (2012) Bioefficacy of novel cyanobacteria-amended formulations in suppressing damping-off disease in tomato seedlings. *World J. Microbiol. Biotechnol.* 28 (12): 3301–3310.
12. Colavita G.M., SperaN., BlackhallV.,& SepulvedaG.M. (2011) Effect of seaweed extract on pear fruit quality and yield. *Acta Hortic.* 909: 601-607.
13. Coppens J.,GrunertO., Van Den Hende S.,VanhoutteI., BoonN., HaesaertG., & De GelderL. (2016) The use of microalgae as a high-value organic slow-release fertilizer results in tomatoes with increased carotenoid and sugar levels. *J. Appl. Phycol.* 28 (4): 2367–2377.
14. DeyB., LernerD.L, LussoP., Boyd M.R.,& ElderJ.H. (2000) Multiple antiviral activities of cyanovirin-N: blocking of human immunodeficiency virus Type 1 gp120 interaction with CD4 and coreceptor and inhibition of diverse enveloped viruses. *J Virol.* 74: 4562-4569 .
15. S. KanagaSubaRaja, C. Viswanathan, (2014) ‘Secured Smart Home Energy Monitoring

- System (SSHEMS) Using RASPBERRY PI', Journal of Theoretical and Applied Information Technology, E-ISSN 1817-3195, Volume 66, Issue 1.
16. Dillehay T.D., Ramirez C., Pino M., Collins M.B., Rossen J., Pino-Navarro J.D. (2008) Monte Verde: seaweed, food, medicine, and the peopling of South America. *Science*. 320, 784–786.
 17. DismukesG.C., CarrieriD., BennetteN., AnanyevG.M.,& PosewitzM.C .(2008) Aquatic phototrophs: efficient alternatives to land-based crops for biofuels. *Curr Opin Biotechnol*. 19: 235–240.
 18. El-ZekyM.M., El-ShahatR.M., MetwalyG.S.,& ElhamM.A. (2005) Using Cyanobacteria or *Azolla* as alternative nitrogen sources for rice production. *J. Agric. Mansoura Univ*. 30(9): 5567-5577.
 19. GantarM., KerbyN.W., RowellP., ObrehtZ., ScrimgeourC.(1995) Colonization of wheat (*Triticum vulgme* L.) by N₂-fixing cyanobacteria: IV. Dark nitrogenase activity and effects of cyanobacteria on natural ¹⁵N abundance in plants. *New Phytol* .129: 337-343.
 20. GuptaV., RathaS.K., SoodA., ChaudharyV.& PrasannaR. (2013) New insights into the biodiversity and applications of cyanobacteria (blue-green algae)—Prospects and challenges. *Algal Res*. 2 (2): 79–97.
 21. Jayachitra S, Prasanth A, (2021) Multi-Feature Analysis for Automated Brain Stroke Classification Using Weighted Gaussian Naïve Baye's Classifier', *Journal of Circuits, Systems, and Computers*, 1-20.
 22. KarthikeyanN., PrasannaR., NainL., & KaushikB.D. (2007) Evaluating the potential of plant growth-promoting cyanobacteria as inoculants for wheat. *Eur. J. Soil Biol*. 43(1): 23–30.
 23. KatirciogluH., TurkeyB.A., BeyatliY., AslimB., YuksekdogZ.,& AticiT. (2006) Screening for antimicrobial agent production of some freshwater. *Microbiology*.2.
 24. KhanW., RayirathU.P., SubramanianS., JitheshM.N., RayorathP., HodgesD.M., etal. (2009) Seaweedextractsasbiostimulantsofplantgrowthand development. *J. Plant Growth Regul*. 28: 386–399.
 25. KongQ.X., LiL., MartinezB., ChenP.,& Ruan,R. (2010) Culture of microalgae*Chlamydomonas reinhardtii* in wastewater for biomass feedstock production. *Applied Biochemistry and Biotechnology*. 160(1), 9.
 26. KudaT., TaniguchiE., NishizawaM., ArakiY. (2002) Fate of water-soluble polysaccharides in

dried *Chorda filum* a brown alga during water washing. J. Food. Comp. Ana. 15: 3 9.

27. KumarM., BauddhK., SaingerM., SaingerP.A., SinghJ.S., & SinghR.P .(2012) Increase in growth,productivity, and nutritional status of rice (*Oryzasativa* L. cv.Basmati) and enrichment in soil fertility applied with an organic matrix entrapped urea. J. CropSci.Biotechnol.15: 137–144.
28. LadhaJ.K.,& ReddyP.M. (2003) Nitrogen fixation in rice systems: state of knowledge and future prospects. Plant Soil, 252: 151-167.
29. LeeR.F.,ValkirsA.O.,& SeligmanP.F. (1989) Importance of microalgae in the biodegradation of tributyltin in estuarine waters. Environ. Sci. Technol. 23: 1515-1518.
30. LeloupM., NicolauR., PallierV., YepremianC.,& Feuillade-CathalifaudG. (2013) Organic matter produced by algae and cyanobacteria: quantitative and qualitative characterization. J. Environ. Sci. 25 (6): 1089–1097.
31. LiX., HuH.Y.,& YangJ. (2010) Lipid accumulation and nutrient removal properties of a newly isolated freshwater microalga,*Scenedesmus* sp.LX1,growing in secondary effluent. N Biotechnol.27(1):59–63.
32. Maurycy Daroch, Shu Geng,& Guangyi Wang. (2013) Recent advances in liquid biofuel production from algal feedstocks,Appl. Energy. 102:1371–1381.
33. MichalakA.M., AndersonE.J., BeletskyD., BolandS., BoschN.S., BridgemanT.B., et al. (2013) Record-setting algal bloom in Lake Erie caused by agricultural and meterological trends consistent with expected future conditions.Proc. Natl. Acad. Sci. USA. 110: 6448-6452.
34. MorandP.,& BriandX. (1996) Excessive growth of macroalgae: a symptom of environmental disturbance. Bot Mar.39: 491-516.
35. MoroneyJ.V.,YnalvezR.A. (2009) Algal photosynthesis. In: eLS, John Wiley & Sons Ltd. Chichester.
36. NoamanN.H., FattahA., Khaleafa M., & ZakyS.H. (2004) Factors affecting antimicrobial activity of *Synechococcus leopoliensis*. Microbiol Res.159:395–402.
37. ObrehtZ., KerbyN.W., GantarM.& RowellP. (1993) Effect of root-associated N₂-fixing cyanobacteria on the growth and nitrogen content of wheat (*Triticum Vulgare*) seedlings. Bio Fert Soils.15: 68-72.

38. OmarH.H. (2000) Nitrogen-fixing abilities of some Cyanobacteria in sandy loam soil and exudates efficiency of rice grain germination. Egypt. J. Phycol.1: 157-167.
39. OswaldW.J., GotaasH.B., GoluekeC.G.,& KellenW.R. (1957) Algae in waste treatment. Sewage Ind Wastes.29:437–55.
40. PattersonG.M., BaldwinC.L., BolisC.M., CaplanF.R., KarusoH., LarsenL.K., et al. (1991) Antineoplastic activity of cultured blue-green algae (Cyanophyta) 1. J Phycol.27:530–6.
41. PlazaM., CifuentesA., & IbanezE. (2008) In the search of new functional food ingredients from algae. Trends Food Sci Tech. 19 (1):31-39.
42. PrasannaR., KanchanA., KaurS., RamakrishnanB., RanjanK., SinghM.C., HasanM., SaxenaA.K.,& ShivayY.S. (2016a) Chrysanthemum growth gains from beneficial microbial interactions and fertility improvements in soil under protected cultivation.Hortic. Plant J. 2 (4): 229–239.
43. PrasannaR., KanchanA., RamakrishnanB., RanjanK., VenkatachalamS., HossainF., ShivayY.S., KrishnanP.,& NainL. (2016b) Cyanobacteria-based bioinoculants influence growth and yields by modulating the microbial communities favorably in the rhizospheres of maize hybrids. Eur. J. Soil Biol. 75: 15–23.
44. PrasannaR., NainL., TripathiR., GuptaV., ChaudharyV., MiddhaS., JoshiM., AnchaR.& Kaushik.B.D. (2008b) Evaluation of fungicidal activity of extracellular filtrates of cyanobacteria– possible role of hydrolytic enzymes. J. Basic Microbiol. 48 (3): 186–194.
45. RathB. (2012) Microalgal bioremediation: Current practices and perspectives. J Biochem Technol.3(3):299–304.
46. Rinehart.K.L., ShawP.D., ShieldL.S., GloerJ.B., HarbourG.C., KohlerM.E.S., et al. (1981) Marine natural products as sources of antiviral, antimicrobial, and antineoplastic agents. Pure Appl. Chem. 53: 795-817.
47. RuffingA.M. (2011) Engineered cyanobacteria: teaching an old bug new tricks. Bioeng. Bugs 2: 136–149.
48. ShobanaS., SarataleG.D., PugazhendhiA., ArvindnarayanS., PeriyasamyS., KumarG.,& KimS.H. (2017) Fermentative hydrogen production from mixed and pure microalgae biomass: Key challenges and possible opportunities. Int. J. Hydrog. Energy. 42(42): 26440-26453.

49. SilkeR., VigdisT., FridaL.D., LiseO.& JorgR. (2007) Nitrogen availability decreases prokaryotic diversity in sandy soils. *Biol. Fertil. Soils.* 43: 449-459.
50. SinghN.K., DharD.W. (2011) Microalgae as second-generation biofuel. A review. *Agron Sustain Dev.*31:605–29.
51. TarakhovshayaE.R., MaslovY.I.,& ShishhovaM.F. (2007) Phytohormones in algae. *Rus. J.PlantPhysiol.* 54, 163–170.
52. ThomasA.D. (2012) Impact of grazing intensity on seasonal variations in soil organic carbon and soil CO₂ efflux in two semiarid grasslands in southern Botswana. *Philos.Trans. R. Soc. Lond. B* 367 (1606): 3076–3086.
53. VerverisC.,& GeorghiouK. (2007) Cellulose, hemicelluloses, lignin and ash content of some organic materials and their suitability for use as paper pulp supplements.*Bioresour. Technol.* 98: 296-30.
54. WangL., LiY., ChenP., MinM., ChenY., ZhuJ.,& RuanR. (2010) Anaerobic digested dairy manure as a nutrient supplement for cultivation of oil-rich green microalgae *Chlorella* sp. *Bioresour Technol.*101(8):2623–8.
55. Wells M.L., Potin P., Craigie J.S., Raven J.A., Merchant S.S., Helliwell K.E., Smith A.G., Camire M.E., Brawley S.H. (2017). Algae as nutritional and functional food sources: revisiting our understanding. *J. Appl. Phycol.* 29, 949–982.
56. Woertz I., FefferA., LundquistT.,& NelsonY. (2009) Algae grown on dairy and municipal wastewater for simultaneous nutrient removal and lipid production for biofuel feedstock.*J Environ Eng.*135(11):1115–22.
57. Yan-GuiS., Xin-RongL., Ying-WuC., Zhi-ShanZ.& YanL. (2013) Carbon fixation of cyanobacterial–algal crusts after desert fixation and its implication to soil organic carbon accumulation in desert. *Land Degrad. Dev.* 24 (4): 342–349.
58. ZhouW., LiY., MinM., HuB., ZhangH., MaX, LiL, ChengY., ChenP.& RuanR. (2012) Growing wastewater-born microalga auxeno *chlorella* *protothecoides* UMN280 on concentrated municipal wastewater for simultaneous nutrient removal and energy feedstock production.*Appl Energy.* 433–440.