

# Estimation and Assessment of Cytogenetic Changes in Bone Marrow Cells of Laboratory Animals Received a Gene-Modified Product

**Nuraliev N.A., Allanazarov A.Kh.**

Bukhara State Medical Institute,  
Urgench branch of the Tashkent Medical Academy

**Key words:** Genetically modified soybeans, cytogenetic changes, bone marrow, polyploidy, aneuploidy, experiment.

The aim was also to study cytogenetic changes in the cells of the bone marrow of laboratory animals that received and did not receive GMO soy. It is found in white mongrel rats treated with GMO soybean detected cytogenetic changes in their red om bone ohm brain e lead partially m the inhibition and proliferation of bone marrow cells. Of all be analyzed metaphases among them revealed altered karyotypes - polyploidy (5.6%) and aneuploidy (5.6%). In the remaining groups (group 2 and 3), no such changes were observed. In the group of animals that did not receive soy with and without GMO, all bone marrow samples had only metaphase plates with a normal karyotype, and the cells did not contain genomic abnormalities.

Genetically modified (GM) organisms are plant or animal organisms, the genotype of which has been changed in a way that is not natural for nature, using methods of genetic engineering to give the body new properties: resistance to herbicides, pests, diseases and salinity, to the action of high and low temperatures, increase calories; to solve the problems of cleaning the environment from organic pollution and heavy metals; to ensure the synthesis of certain compounds in the plant organism and the use of plants for the production of these compounds [ 3, 9, 10 ].

Genetic changes are made for scientific or economic purposes and distinguished by a purposeful change in the genotype of an organism, in contrast to the random, characteristic of a natural mutation process. The main trait inherent in most GMO plants is herbicide and pest resistance. Cultivation of GM-soybeans does not have a single positive moment [ 5, 7, 12].

The most serious risks associated with products of genetic engineering are combined into 2 main groups: food and environmental [ 2, 8].

Nutritional - weakening of the immune system; the occurrence of allergic reactions; the development of pathologies associated with the accumulation of pesticides in the human body, introduced by the used GMO products, including GMOsoy.

Ecological - the loss of the diversity of the gene pool of wild relatives of cultivated plants due to their cross-pollination with related GM plants; o contamination of water resources by the use of pesticides; depletion and disturbance of natural soil fertility associated with the suppression of GM plants by toxins, the vital activity of soil invertebrates and microflora; Acceleration of the second plant growth to a much greater extent than in traditional agricultural RMS households affiliations cultures. In general, the above studies indicate the insufficient substantiation of the safety of GMO plant products for humans and the environment.

Current trends in the change in the structure of morbidity indicate an increase in the relative importance of genetically determined diseases [ 8, 9].

Many authors believe that now important is to solve the urgent problems and correlating the rate of changes in the environment with the ability to adapt human populations, in addition it is necessary to assess the extent to which, in the biosphere changes that affect the rate of mutation [ 6, 7, 11].

The results of biological monitoring of human populations have shown that at present there is not only an increase in hereditary pathology, but also an increase in the frequency of diseases with a genetic component. The danger of induced mutagenesis is that newly emerging mutations have a negative impact on the fitness of the population and on the health of the population.

In this regard, new research in this direction is relevant and in demand at the present time.

**The aim of the study** was to learn the evaluation cytogenetic changes in bone marrow cells in laboratory animals, treated and not treated with GMO-product in a comparative aspect.

**Materials and methods.** To perform the planned research used 90 white outbred rats weighing bodies 150-180 g of both sexes, contained in a standardized vivarium conditions- with relative humidity 50-60%, temperature vivarium 22<sup>0</sup>C, light conditions (12 h aces darkness and light). The maintenance of laboratory animals, feeding and caring for them, selection of animals, cleaning and disinfection of the vivarium premises were carried out in accordance with the approved recommendations [ 4]. All laboratory animals were obtained from the same nursery and were of the same age. All laboratory animals were quarantined for 21 days prior to the start of experimental studies.

When working with experimental animals, all ethical principles of working with laboratory animals and the rules of biological safety were strictly observed [4].

When conducting cytogenetic studies, all operations when working with growth media and preparations were performed under sterile conditions using a laminar box. Buffers were prepared with double-distilled water, filtered through membrane filters (0.22 um « Millipor », Germany) and autoclaved at 1.2 bar for 30 minutes. Before use, glassware is pre-sterilized at 160<sup>0</sup>C for 120 minutes. Equipment, fixtures, utensils made of polymeric materials were exposed to ultraviolet light for 30 minutes.

For experimental studies, the bone marrow was collected from the femur of laboratory animals during dissection of the animal.

All laboratory animals were divided into the following groups:

1 - group - white outbred rats (n = 12), receiving GMO soy with food for 30 days at 0.02-0.03 grams per one laboratory animal;

Group 2 - non-bred white rats (n = 12), treated without GMO-modified soy;

Group 3 - non-pedigree white rats (n = 6), who received a standard diet without GMO soy and conventional soy.

As a GMO product in experiments used to I grown overseas and imported into our country only to perform scientific research of their work (see Fig. 1).



**Figure: 1. Appearance of GMO soybeans**

Using the method of polymerase chain reaction (PCR), the presence of the 35 S + FMV promoter in the studied GMO- soy was revealed , which proves that the studied soy is a GMO-product.

Cytogenetic changes in rat bone marrow cells were studied using the following methods:

Direct method. Red bone marrow was washed out of the femur with a nutrient medium with 0.04% colchicine into a centrifuge tube and incubated for 2-2.5 hours in a thermostat at 37 °C. Then it was further incubated with a hypotonic solution of KS L for 40 minutes in a thermostat at 37 °C. After hypotonization, it is treated three times with a fixative in the proportion of one part of glacial acetic acid and three parts of 960-1000 ethyl alcohol. The resulting precipitate was applied to a previously cleaned defatted glass slide and stained with Giemsa dye.

Method according to Ford. After the end of the drug administration, all the animals were injected intraperitoneally with 0.5 ml of a 0.1% colchicine solution, after 2 hours the animals were killed in strict accordance with the ethical principles of working with laboratory animals. Red bone marrow flushed from a femur hypotonic solution COP L and incubated for 40 minutes in an oven at 37 °C. After hypotension, they were treated three times with a fixative in the proportion of one part of glacial acetic acid and three parts of 960-1000 ethyl alcohol. The resulting precipitate was applied to a previously cleaned defatted glass slide and stained with Giemsa dye.

Statistical processing was carried out by the generally accepted methods of variation statistics.

**Research results.** For the analysis, cells of the red bone marrow were used, in which elements of the mitotic apparatus were detected. Metaphase plates - this cells in which chromosomes are in the metaphase mitotic (somatic cell division stage). The number of chromosomes in rats is normally 42 (diploid set).


Analysis revealed that - the low mitotic activity of bone marrow cells in 1 - group (treated with GMO -soy) only 3/10 animals in the bone marrow samples detected metaphase plate (tab.).

**Table**  
**Indicators of cytogenetic changes in bone marrow cells of laboratory animals treated with a GMO product (soy)**


Group	Types of aberrations		
	Polyploidy	Aneuploidy	Structural changes in chromosomes
1 - group	5,6%	5,6%	Dispersion, Pulverization, Delayed mitosis in prophase
2 - group	Lack of	Lack of	Pulverization of chromosomes, K-mitosis
3 - group	Lack of	Lack of	Lack of

Of all the metaphases to be analyzed in laboratory animals of the 1st group, in 11.2% cases, polyploid (the cell contains a set of chromosomes more by a multiple than normal; 5.6%) and aneuploid (cells contain the number of chromosomes less or more than a multiple haploid set; 5.6%) karyotypes.


In other bone marrow samples from rats of this experimental group, metaphase plates were absent, but there were blast cells (cells at the stage preceding mitosis) and cells with mitotic pathology: chromosome pulverization (Fig. 2 and Fig. 3), impaired chromosome spiralization and despiralization (Fig. 4), premature spiralization of chromosomes (Fig. 5), delayed mitosis at the prophase stage. The rest of the metaphase plates contained a normal (42 chromosomes) karyotype.




**Figure: 2. Bone marrow cells of laboratory animals treated with GM-soy (group 1). Pulverization of chromosomes (approx. X 10, vol. X 100)**



**Figure: 3. Bone marrow cells of laboratory animals treated with GM-soy (group 1). Pulverization of chromosomes (approx. X 10, vol. X 100)**



**Figure: 4. Bone marrow cells of laboratory animals treated with GM-soy (group 1). Violation of spiralization and despiralization of chromosomes (approx. X 10, Ob. X 100)**




**Figure: 5. Bone marrow cells of laboratory animals treated with GM-soy (group 1).  
Premature spiralization of chromosomes  
(Approx. X 10, Ob. x 100)**


Experimental studies on laboratory animals proved that the used GM product (soy) led to cytogenetic changes in the cells of highly proliferating tissues, such as the red bone marrow of white outbred rats.

There is a partial inhibition of the proliferation of bone marrow cells due to the cytotoxic action, which can lead to cytopenia.

In group 2, polyploid (polyploidy) and aneuploid (aneuploidy) karyotypes were not detected in only 50% of laboratory animals. The number of metaphase plates found in the bone marrow was the same with the animals of group 1. In all analyzed metaphases, the karyotype was not changed (Fig. 6 and Fig. 7).



**Figure: 6. Bone marrow cells of laboratory animals that received soy without GM (group 2).  
Normal karyotype (approx. X 10, vol. X 100)**




**Figure 7. Bone marrow cells of laboratory animals that received soy without GM (group 2). Normal karyotype (approx. X 10, vol. X 100)**

It was found that the cells did not contain genomic abnormalities, all metaphases had a normal karyotype (42 chromosomes). In the remaining (50%) bone marrow samples, cells with mitotic pathology (chromosome pulverization, K-mitosis) were present, as well as in the bone marrow cells of group 1 outbred white rats.

In group 3 (control group), all bone marrow samples contained only metaphase plates with a normal karyotype (Fig. 8 - early metaphase; Fig. 9 - late metaphase).



**Figure 8. Bone marrow cells of laboratory animals that received a standard diet (group 3 - control). Normal karyotype, early metaphase. (Approx. X 10, Ob. x 100)**



**Figure: 9. Bone marrow cells of laboratory animals fed a standard diet (group 3 - control).  
Normal karyotype, late metaphase (approx. X 10, vol. X 100)**

The mitotic activity of rat bone marrow cells was higher than in the experimental groups (groups 1 and 2).

It has been established that to establish the true mutagenesis in the chromosomes of cells, it is necessary to test and investigate cytogenetic changes with a single exposure to the first mitoses (24 hours after exposure).

### **Conclusions.**

1. It was found that in the group of white outbred rats receiving GM-soy, cytogenetic changes were found in their red bone marrow, expressed in partial inhibition of bone marrow cell proliferation due to cytotoxic action, which can lead to cytopenia.

2. Of all the metaphases to be analyzed in laboratory animals of the 1st group, in 11.2% of cases, altered karyotypes were revealed - polyploidy (5.6%) and aneuploidy (5.6%). In the remaining groups (group 2 and 3), no such changes were observed.

3. Bone marrow samples from 1-group rats contained blast cells and cells with mitotic pathology: chromosome pulverization, chromosome spiralization and despiralization disorders, premature chromosome spiralization, mitosis delay at the prophase stage.

4. In the group of laboratory animals as feed given to w without GM (group 2) was found to decrease the proliferation of bone marrow cells (at 50%), the rest in the bone marrow were present metaphase plate in all metaphases analyzed (36) the karyotype was not changed. The cells did not contain genomic abnormalities; all metaphases had a normal karyotype (42 chromosomes).

5. In the group of animals that did not receive soy with and without GM (3 - groups a control group), all bone marrow samples contained only metaphase plates with normal karyotype, and the cells did not contain genomic abnormalities.

### **References**

1. Derevenschikova M.I., Syromyatnikov M.Yu. The use of molecular genetic methods for microbiological control of food products // Technics and technology of food production . - Moscow, 2018 . - No. 4. - P.87-113.

2. Dronik G.V., Black I.V. The content of middle molecules in the blood serum of the rats in the use of conventional and transgenic soybean // Bulletin of the present-day e variables studies. - Moscow, 2017. - No. 6-1. - With .7-8.
3. Zakirova G.Sh., Papunidi K.Kh., Kadikov I.R. The effect of diets with the content of genetically modified soybeans in the body of the animal 's // veterinarian. - Moscow, 2019. - No. 4. - S. 37-43.
4. Nuraliev N.A., Bektimirov A.M.-T., Alimova M.T., Suvonov K.Zh. Rules and methods of working with laboratory animals in experimental microbiological and immunological research // Methodical manual. - Tashkent, 2016.-- 33 p.
5. Pavlovskaya N.E., Lushnikov A.V., Polekhina N.N., Solokhina I.Yu., Gneusheva I.A. Influence of genetically modified soybeans on body weight and reproductive activity of white laboratory mice // Bulletin of Science and Education. - Moscow, 2018. - No. 2. - With .1-12.
6. Sobirova D.R., Nuraliev N.A., Ginatullina E.N. Results of experimental studies on the study and assessment of the mutagenic activity of a genetically modified product // Problems of Biology and Medicine. - Samarkand, 2017. - No. 1 . - S. 180-184.
7. Sobirova D.R., Nuraliev N.A., Nosirova A.R., Ginatullina E.N. Study of the effect of a genetically modified product on mammalian reproduction in experiments on laboratory animals // Infection, immunity and pharmacology. - Tashkent, 2017. - No. 2. - S. 195-200.
8. Tutelyan V.A. Ensuring the safety of genetically modified organisms for food production // Herald of the Russian A kademii N AUC. - Moscow, 2017.-- Volume 87 . - No. 4. - S.342-347.
9. Tyshko N.V. Control over genetically modified organisms of plant origin in food products: scientific substantiation and methodological support // Questions of nutrition. - Moscow, 2017.-- Volume 86 . - No. 5. - pp. 29-33.
10. Global Status of Commercialized Biotech / GM Crops in 2014, Pocket K No.16, ISAAA, p. 5 .
11. Nuraliev NA, Sobirova DR, Baltaeva KA, Ginnatullina EN Effect of genetically modified product on reproduction function, biochemical and hematology indexes in experimental study // European Science Review. - Austria, Vienna, 2017.-- N1. - P.94-95.
12. Willems S. Fraiture M.A. , Deforce D. , De Keersmaecker S.C. , De Loose M. , Ruttink T. , Herman P. , Van Nieuwerburgh F. , Roosens N. Statistical framework for detection of genetically modified organisms based on Next Generation Sequencing // Food Chemistry. - 2016. - Vol. 192. - P.788-798.
13. Bazarova D. Some problems of counteracting crimes related to laundering of illegal proceeds in Uzbekistan Journal of Advanced Research in Dynamical and Control Systems. Volume 11, Issue 7, 2019, Pages 873-885
14. Ismailova, Z., Choriev, R., Ibragimova, G., Abdurakhmanova, S., & Abdiev, N. (2020). Competent model of Practice-oriented education of students of the construction profile. Journal of Critical Reviews. Innovare Academics Sciences Pvt. Ltd. <https://doi.org/10.31838/jcr.07.04.85>
15. Ismailova, Z., Choriev, R., Musurmanova, A., & Aripjanova, M. (2020). Methods of training of teachers of university on advanced training courses. Journal of Critical Reviews. Innovare Academics Sciences Pvt. Ltd. <https://doi.org/10.31838/jcr.07.05.85>
16. Ismailova, Z., Choriev, R., Salomova, R., & Jumanazarova, Z. (2020). Use of economic and geographical methods of agricultural development. Journal of Critical Reviews. Innovare Academics Sciences Pvt. Ltd. <https://doi.org/10.31838/jcr.07.05.84>

17. Isakov, A., Tukhtamishev, B., & Choriev, R. (2020). Method for calculating and evaluating the total energy capacity of cotton fiber. IOP Conference Series: Earth and Environmental Science, 614(1), 012006
18. Davirov, A., Tursunov, O., Kodirov, D., Baratov, D., & Tursunov, A. (2020). Criteria for the existence of established modes of power systems. IOP Conference Series: Earth and Environmental Science, 2020, 614(1), 012039
19. Obidov, B., Choriev, R., Vokhidov, O., & Rajabov, M. (2020). Experimental studies of horizontal flow effects in the presence of cavitation on erosion-free dampers. IOP Conference Series: Materials Science and Engineering, 883(1), 012051
20. Khasanov, B., Choriev, R., Vatin, N., & Mirzaev, T. (2020). The extraction of the water-air phase through a single filtration hole. IOP Conference Series: Materials Science and Engineering, 2020, 883(1), 012206
21. Shokhrud F. Fayziev The problem of social stigma during a pandemic caused by COVID-19 International Journal of Advanced Science and Technology Vol. 29, No. 7, (2020), pp. 660-664 <http://sersc.org/journals/index.php/IJAST/article/view/13965/7188>
22. Fayziyev Shokhrud Farmonovich Medical law and features of legal relations arising in the provision of medical services. International journal of pharmaceutical research Volume 11, Issue 3, July - Sept, 2019 P. 1197-1200 doi:10.31838/ijpr/2019.11.03.088 <http://www.ijpronline.com/ViewArticleDetail.aspx?ID=11016>
23. Bryanskaya Elena, Fayziev Shokhrud, Altunina Anna, Matiukha Alena Topical Issues of an Expert Report in the Process of Proving in a Criminal Examination. International Journal of Engineering and Advanced Technology (IJEAT) ISSN: 2249 – 8958, Volume-9 Issue-1, October 2019 5345-5349 DOI: 10.35940/ijeat.A2946.109119 <https://www.ijeat.org/wp-content/uploads/papers/v9i1/A2946109119.pdf>
24. Fayziev Shokhrud (2019) Legal Aspects of Transplantology in the Republic of Uzbekistan. Systematic Reviews in Pharmacy, ISSN: 0976-2779, Vol: 10, Issue: 2, Page: 44-47 doi:10.5530/srp.2019.2.08 <http://www.sysrevpharm.org//fulltext/196-1575419211.pdf?1586863081>
25. Tulaganova, G. Some issues of observance of international legal norms of fight against legalization of criminal incomes in the Republic of Uzbekistan Journal of Advanced Research in Dynamical and Control Systems 12(2 Special Issue), c. 143-155
26. Bekchanov D; Kawakita H; Mukhamediev M; Khushvaktov S; Juraev M. Sorption of cobalt (II) and chromium (III) ions to nitrogen- and sulfur- containing polyampholyte on the basis of polyvinylchloride /Polymers for Advanced Technologies 2021 <https://doi.org/10.1002/pat.5209>
27. Davron, B., Mukhtar, M., Nurbek, K., Suyun, X., Murod, J. Synthesis of a New Granulated Polyampholyte and its Sorption Properties. *International Journal of Technology*. Volume 11(4), pp. 794-803. ., (2020) <https://doi.org/10.14716/ijtech.v11i4.4024>
28. Mukhamediev, M.G., Bekchanov, D.Z. New Anion Exchanger Based on Polyvinyl Chloride and Its Application in Industrial Water Treatment. *Russ J Appl Chem* 92, 1499–1505 (2019). <https://doi.org/10.1134/S1070427219110053>
29. Mukhamediev, M.G., Auelbekov, S.A., Sharipova, Z.T. et al. Polymer complexes of gossypol and their antiviral activity. *Pharm Chem J* 20, 276–278 (1986). <https://doi.org/10.1007/BF00758817>
30. Ikramova, M.E., Mukhamediev, M.G., Musaev, U.N. Complexation of hydrazine- and phenylhydrazine-modified nitron fibers with iodine/ *Plasticheskie Massy: Sintez Svoystva Pererabotka Primenenie*, (12), crp. 41–45 (2004)

31. Gafurova, D.A., Khakimzhanov, B.S., Mukhamediev, M.G., Musaev, U.N. Sorption of Cr(VI) on the anion-exchange fibrous material based on nitron. Russian Journal of Applied Chemistry, 75(1), стр. 71–74, (2002)

32. Rustamov, M.K., Gafurova, D.A., Karimov, M.M. et al. Application of ion-exchange materials with high specific surface area for solving environmental problems. Russ J Gen Chem 84, 2545–2551 (2014). <https://doi.org/10.1134/S1070363214130106>