

Are We Comprehensively Treating Patients From Assessment To Completion Of Treatment Plan?- A Retrospective Analysis

Type of manuscript: Research article

Running title: Assessment to completion of comprehensive treatment plan

Ranjith Raj VPRB

Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical sciences (SIMATS)
Saveetha University,
Chennai - 600077
E-Mail ID: 151701035.sdc@saveetha.com
Contact:+919444727283

GifrinaJayaraj

Reader,
Department of Oral Pathology
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical sciences (SIMATS)
Saveetha University,
Chennai - 600077
Email ID: gifrina@saveetha.com
Contact Number:+919952096111

MuruganThamaraiselvan

Reader
Department of Periodontics,
Saveetha Dental College,
Saveetha Institute of Medical and Technical sciences (SIMATS)
Saveetha University, Chennai - 600077
E-Mail ID: thamaraiselvan@saveetha.com
Contact:+918610507379

Corresponding Author

GifrinaJayaraj

Reader,
Department of Oral Pathology
Saveetha Dental College and Hospitals,
Saveetha Institute of Medical and Technical sciences (SIMATS)

Saveetha University, Chennai - 600077

Email ID:gifrina@saveetha.com

Contact Number:+919952096111

Total No: Of Words:4897

Abstract

Aim:

Any treatment is designed and performed to increase the quality of life of the patient. The success of any treatment depends on how far the patient is relieved from the disease. This will ultimately lead to the improvement in the health and well being of the patient. The formulation, the execution and the completion of a treatment plan requires the complete determination from the patient as well as the practitioner and this is dependent upon various factors. This study aims comprehensively analyse the assessment of the patient till the completion of treatment plan in a dental institution.

Materials And Methods:

A retrospective cross sectional study was done by assessing 86000 patient records who visited as an outpatient in a dental institution in south India from June 2019 to March 2020. Data of 100 random patients were obtained by random sampling to prevent bias. Fields such as unique patient ID, treatment completed, treatment under progress and total treatment planned, type of case: emergency/normal, Type of Treatment plan: Interim/Cumulative were recorded in Microsoft Excel Software and then exported to IBM SPSS Statistics Software for further analysis where frequency, chi-Square & descriptive statistics were done.

Results:

Average percentage of treatment completed 60.05%(SD 66.67), Number Of emergency cases :33, Number Of Normal cases:67, average percentage of treatment pending 39.95%(SD=33.33), Total number of treatment planned ranged from 1-31, 100 % of the treatment was completed majorly in cases with less than 5 treatment procedures(20%). At least one planned treatment was completed for most of the patients.(88%)

Conclusion:

The process of diagnosis, planning and execution of a treatment plan for each patient is unique and significant usually in emergency cases and cases with aesthetic requirements, the patients are determined for relieving the pain & resolving their aesthetic needs and are not willing to complete the treatment plan. Hence proper knowledge, awareness and practice is required for both the patient as well as the practitioners.

Keywords: Comprehensive, patients, prognosis, treatment, treatment plan.

Introduction

Oral health of a patients is one of the significant factor attributing to their overall health and well being.[1–3] Maintaining one's oral health is in the hands of the patients themselves as well as the practitioner.[4–6] It is the duty of the dental practitioner to completely assess , diagnose the patient's needs, formulate ,execute and complete the treatment plan for a patient. [7–9] Any treatment is designed and performed to increase the quality of life of the patient.[10–12] The success of a treatment depends on how far the patient is relieved from the disease.This will ultimately lead to the improvement in the health and overall well being of the patient.[6,13,14]

Dental practitioners are responsible for formulating the treatment plan. The clinician evaluates the oral cavity for the quality of its health. This requires years of theoretical,practical and clinical knowledge.[15–17] The evaluation is done with the help of various diagnostic tools, which is categorised into radiographics, non radiographic and there are many recent advances aiding in the proper diagnosis. [18–20]

The proper diagnosis involves detailed history taking, which is relevant to the patient's needs. [21–23] The formulation and completion of treatment plan involves various factors involving the patient as well as the clinician such as systemic health, financial capabilities, preference of the patients, knowledge of the dentist, experience of the practitioner , their training in their own field, laboratory support, functional , aesthetic and technical demands.[24–26] Treatment plan to each patient is unique just as a fingerprint. [24–26]

Comprehensive treatment planning involves resolving the chief complaint followed by regular recall and follow ups to complete the other disease manifestations the patient is unaware of. There are various methods in assessment,evaluation and formulating treatment plan followed in various dental schools all around the world (tokede O et al.) did the first investigation to quantitatively report on student treatment planning performance in dentistry.[27–29] There has been limited studies on treatment planning,diagnosis and the outcomes of the treatment individually.Our team has rich experience in research and we have collaborated with numerous authors over various topics in the past decade [30–54]. This study aims to comprehensively study the completion of treatment plans,identify reasons for incomplete treatments,and overcome hurdles regarding the same.

Materials And Methods

Study Design And Study Setting

This retrospective cross-sectional study analysed the records of outpatients, visiting saveetha dental college,chennai. Ethical approval was obtained from the institutional ethical review boards. The study population included 100 patients by random sampling in order to prevent bias.

Data Collection

Data was obtained assessing 86000 patient records who visited as an outpatient in a dental institution in south India from june 2019 to march 2020. Data of 100 random patients were

obtained by random sampling to prevent bias. Parameters such as Total treatment planned, Number of Treatments completed, Treatment under progress/pending, Type of case-Normal/emergency, Type of treatment Plan:Interim/cumulative were noted and recorded in Microsoft Excel 2004.

Statistical Analysis

The data recorded in Microsoft Excel was exported to IBM Corp. Released 2011. IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp. Software Descriptive statistics-frequencies, Chi-square tests were performed. $p < 0.05$ was considered statistically significant.

Results

A Total of 100 patients' treatment plan completion was included for analysis. Among which Emergency cases were 33% and Normal cases were 67%. [Figure :1] Total Number of treatments planned ranges from 1-31 procedures. Majority of the study population had less than 5 treatment procedures planned (56%) followed by 6-10 procedures (29%), 11-15 procedures (9%), 16-20 procedures (3%) and more than 20 procedures was only by (3%). Emergency cases had less number of treatment procedures planned compared to normal visit patients. However, this difference was found to be statistically not significant ($p = 0.117$) [Figure:2,3]

Average percentage of treatment completed 66.91% (SD=72.73). [Table:2] At least one planned treatment was completed for most of the patients (88%). No treatment was initiated for 12% of the study population, 26% had completed their comprehensive treatment plan. [Figure:4]. Percentage of treatment completed in emergency cases (33%) was comparatively lesser than percentage of treatment completed in normal cases (67%). However, this difference was found to be statistically not significant ($p = 0.273$) [Figure:5] 100 % of the treatment was completed majorly in cases with less than 5 treatment procedures (20%). The more number of procedures in the treatment plan was inversely related to the percentage of treatment completion. This difference was however not statistically significant ($p = 0.099$) [Figure:6]

Discussion

A treatment plan is a carefully planned out sequence of treatments designed to eliminate the etiological factor of disease. A comprehensive treatment is built by proper history taking with logical progression to the diagnosis of the disease. [55] Comprehensive treatment plan is defined as a treatment plan which satisfies the main four domains such as the function, symptomatology, pathology, aesthetics. [56] Function refers to the restoration of the function to the maximum extent, but no restoration can restore the complete functioning of a natural tooth. [57] Symptomatology refers to the symptoms exhibited by the patient and the complete relief from the symptoms. [58] Pathology refers to resolving the main root cause of the disease and analysing how it progressed. [59] Aesthetics which is often not taken into concern in general

medicinal treatment, dentistry can be compared to that of surgery but aesthetics is of much importance in dentistry as, the teeth remain the most noted aspect of an individual's face and must be aligned and restored according to the patient's wishes as aesthetics is subjective to each and every individual.[60,61] (Sorin T et al.) state that in a comprehensive approach to a individual patient, The profile of the patient should be updated & provided regularly. This requires frequent recalls and reviews.[62]

The need for a comprehensive to any dental ailment is high because, treating the symptoms such as pain in a short term will only be a temporary relief and has high chances of recurrence, hence proper diagnosis and the analysis of the root cause of the ailment is required to completely eradicate the problems faced by the patients.[63]

In this retrospective study it is reported that 67% of the cases are normal elective procedures, and 33 % of the cases are emergency procedures.[Figure:1] The main motive of emergency procedure is to relieve the pain and other symptoms in the short term.[64]

The total number of treatments ranges from 1-31 and it can vary from person to person depending upon the oral health [Figure:2,3], their habits and various other factors. Average percentage of treatment completed is 66.91% (SD=72.73) [Table:2] . It is appreciable, however 100% treatment completion will be most appreciated, steps should be taken to attain it and will ultimately be attributed to the patient as well as the practicing clinician.

Average percentage of treatment pending 45.09% (SD=65.89) [Table:2] and no treatment has been done for 12% of the study population [Figure:4], percentage of treatment completed in emergency cases (33%) was comparatively lesser than percentage of treatment completed in normal cases (67%) [Figure:5] the reason for this could be due to emotional factors, financial factors, willingness of the patient, availability of the doctor, availability of the lab equipment.

A treatment or a treatment plan could fail due to various reasons clinical errors such as improper diagnosis, errors in the interpretation of the lab investigations, iatrogenic errors.[65] The success of a treatment and a treatment plan not only involves on the clinician and their skills, but also on the patient, it is the duty of the patient to immediately aid the help of a dental practitioner as soon as they start to feel any difficulty in their oro facial region, the more the patient delays to consult a dental practitioner the severity of the ailment will increase will be difficult to the patient and as well as the practitioner.[66] Other psychological factors such as dental anxiety and fear are also some of the factors resulting in the delaying of a dental treatment.[67] These psychological factors can be reduced by conducting campaigns in schools, awareness programmes by the local dental association can also be conducted to raise the knowledge and awareness among the general population.[65,68]

It also the duty of the patient to coordinate with their practitioner to abide to the treatments, the prognosis of a treatment is mainly dependent on the patient to properly maintain the restoration, oral health and oral hygiene. In a study done by (Mettes D) Insufficient data, inability to recall patients, improper follow ups were some of the difficulties faced in the completion of a treatment plan.[69]

Attitude of the patient and the dental practitioner towards the treatment also plays a major role in a successful treatment, this can be improved by having a well communicable interactional relationship between the patient, so that both the patient and as well as the doctor can share their difficulties, ideas in terms of the treatment which is to be or had been executed. 100 % of the treatment was completed majorly in cases with less than 5 treatment procedures (20%) .[Figure:6] A proper communication can also help in planning the treatment, appointments and recalls ultimately leading to the success of a treatment.[70] (Patel S et al.) concluded that there is not enough evidence to conclude that one type of recall protocol can be followed for every patient. Each treatment plan should be regarded and treated uniquely.[71,72] Despite the limitation of the previous studies, our study serves as the first study to analyse a comprehensive treatment plan in an institution. Our study has few limitations such as a smaller sample has only been analysed, financial & emotional status of the patient was not taken into account. General awareness not known to the specific cohort. Our institution is passionate about high quality evidence based research and has excelled in various fields ([16,73–82] This study serves as a pilot study and further studies may be carried out with a larger population, better control groups and include other variables like emotion, education, and patient motivation.

Conclusion

It is evident that completion of the treatment plan is essential for both the practicing clinicians as well as the patient getting the treatment. The process of diagnosis, planning and execution of a treatment plan for each patient is unique and significant, usually in emergency cases and cases with aesthetic requirements, the patients are determined for relieving the pain, symptoms & resolving their aesthetics needs and are not willing to complete the treatment plan. Also, more number of planned procedures result in poor patient compliance and inability to complete treatment procedures. Hence proper knowledge awareness and practice is required for both patients as well as practitioners to support the patients and reinforce the importance of completion of planned treatments for the benefit of the patient.

Acknowledgment

Nil.

Conflict of interest

The authors declare no conflict of interest.

Authors contribution

Author 1(Ranjith Raj VPRB) carried out the retrospective study by collecting the data and prepared the draft of the manuscript after performing the required statistical analysis. Author 2(Gifrina Jayaraj) helped in the conception of the topic, and contributed in the study design, statistical analysis and aided in the drafting of the manuscript. Author

3(MuruganThamaraiselvan) contributed in developing and formatting the manuscript.All the above mentioned authors have discussed among themselves and have contributed to this study.

References

1. AL-Omiri MK, Karasneh J. Relationship between Oral Health-Related Quality of Life, Satisfaction, and Personality in Patients with Prosthetic Rehabilitations [Internet]. Vol. 19, Journal of Prosthodontics. 2010. p. 2–9. Available from: <http://dx.doi.org/10.1111/j.1532-849x.2009.00518.x>
2. Sridharan G, Ramani P, Patankar S. Serum metabolomics in oral leukoplakia and oral squamous cell carcinoma. J Cancer Res Ther. 2017 Jul;13(3):556–61.
3. Thangaraj SV, Shyamsundar V, Krishnamurthy A, Ramani P, Ganesan K, Muthuswami M, et al. Molecular Portrait of Oral Tongue Squamous Cell Carcinoma Shown by Integrative Meta-Analysis of Expression Profiles with Validations. PLoS One. 2016 Jun 9;11(6):e0156582.
4. Naito M, Yuasa H, Nomura Y, Nakayama T, Hamajima N, Hanada N. Oral health status and health-related quality of life: a systematic review. J Oral Sci. 2006 Mar;48(1):1–7.
5. Gupta V, Ramani P. Histologic and immunohistochemical evaluation of mirror image biopsies in oral squamous cell carcinoma. Journal of Oral Biology and Craniofacial Research. 2016 Sep 1;6(3):194–7.
6. Jayaraj G, Sherlin HJ, Ramani P, Premkumar P, Natesan A. Stromal myofibroblasts in oral squamous cell carcinoma and potentially malignant disorders. Indian J Cancer. 2015 Jan;52(1):87–92.
7. House MM. The relationship of oral examination to dental diagnosis [Internet]. Vol. 8, The Journal of Prosthetic Dentistry. 1958. p. 208–19. Available from: [http://dx.doi.org/10.1016/0022-3913\(58\)90150-1](http://dx.doi.org/10.1016/0022-3913(58)90150-1)
8. Sivaramakrishnan SM, Ramani P. Study on the Prevalence of Eruption Status of Third Molars in South Indian Population. Biol Med [Internet]. 2015;07(04). Available from: <https://www.omicsonline.com/open-access/study-on-the-prevalence-of-eruption-status-of-third-molars-in-south-indian-population-0974-8369-1000245.php?aid=59465>
9. Shree KH, Hema Shree K, Ramani P, Herald Sherlin, Sukumaran G, Jeyaraj G, et al. Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma – a Systematic Review with Meta Analysis [Internet]. Vol. 25, Pathology & Oncology Research. 2019. p. 447–53. Available from: <http://dx.doi.org/10.1007/s12253-019-00588-2>

10. Viveka TS, Shyamsundar V, Krishnamurthy A, Ramani P, Ramshankar V. p53 Expression Helps Identify High Risk Oral Tongue Pre- malignant Lesions and Correlates with Patterns of Invasive Tumour Front and Tumour Depth in Oral Tongue Squamous Cell Carcinoma Cases. *Asian Pac J Cancer Prev.* 2016;17(1):189–95.
11. Hannah R, Ramani P, Herald. J. Sherlin, Ranjith G, Ramasubramanian A, Jayaraj G, et al. Awareness about the use, Ethics and Scope of Dental Photography among Undergraduate Dental Students Dentist Behind the lens. *Research Journal of Pharmacy and Technology.* 2018 Mar 30;11(3):1012–6.
12. Sischo L, Broder HL. Oral health-related quality of life: what, why, how, and future implications. *J Dent Res.* 2011 Nov;90(11):1264–70.
13. Jangid K, Alexander AJ, Jayakumar ND, Varghese S, Ramani P. Ankyloglossia with cleft lip: A rare case report. *J Indian Soc Periodontol.* 2015 Nov;19(6):690–3.
14. Iqbal MK, Kim S. A review of factors influencing treatment planning decisions of single-tooth implants versus preserving natural teeth with nonsurgical endodontic therapy. *J Endod.* 2008 May;34(5):519–29.
15. Wyrwich KW, Bullinger M, Aaronson N, Hays RD, Patrick DL, Symonds T, et al. Estimating clinically significant differences in quality of life outcomes. *Qual Life Res.* 2005 Mar;14(2):285–95.
16. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med.* 2019 Apr;48(4):299–306.
17. Swathy S, Gheena S, Varsha SL. Prevalence of pulp stones in patients with history of cardiac diseases. *Research Journal of Pharmacy and Technology.* 2015;8(12):1625–8.
18. Sundaram N, Narendar R, Dineshkumar P, Ramesh S, Gokulanathan S. Evaluation of oral health related quality of life in patient with mild periodontitis among young male population of Namakkal district [Internet]. Vol. 5, *Journal of Pharmacy and Bioallied Sciences.* 2013. p. 30. Available from: <http://dx.doi.org/10.4103/0975-7406.113289>
19. Kalenderian E, Tokede B, Ramoni R, Khan M, Kimmes N, White J, et al. Dental clinical research: an illustration of the value of standardized diagnostic terms. *J Public Health Dent.* 2016 Mar;76(2):152–6.
20. Huang K-C, Lai C-H, Huang C-F, Lu H-K. A comprehensive periodontal treatment project: The periodontal status, compliance rates, and risk factors. *J Dent Sci.* 2016 Jun;11(2):182–

8.

21. De Franco RL. Essentials of complete denture Prosthodontics. 1988;384–482.
22. Kumar A, Sherlin HJ, Ramani P, Natesan A, Premkumar P, Others. Expression of CD 68, CD 45 and human leukocyte antigen-DR in central and peripheral giant cell granuloma, giant cell tumor of long bones, and tuberculous granuloma: An immunohistochemical study. *Indian J Dent Res.* 2015;26(3):295.
23. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species–mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol* [Internet]. 2019; Available from: <https://journals.sagepub.com/doi/abs/10.1177/0960327119839173>
24. Sivakumar A, Thangaswamy V, Ravi V. Treatment planning in conservative dentistry. *J Pharm Bioallied Sci.* 2012 Aug;4(Suppl 2):S406–9.
25. Henderson D, McCracken WL, McGivney GP, Castleberry DJ. McCracken’s removable partial prosthodontics. CV Mosby Company; 1985.
26. Carranza FA, Newman M. Clinical Periodontology. New York. WB Saunders; 1996.
27. Fasbinder DJ. Treatment planner’s toolkit. *Gen Dent.* 1999 Jan;47(1):35–9.
28. Tokede O, Walji M, Ramoni R, White JM, Schoonheim-Klein M, Kimmes NS, et al. Treatment planning in dentistry using an electronic health record: implications for undergraduate education. *Eur J Dent Educ.* 2013 Feb;17(1):e34–43.
29. Jayaraj G, Sherlin HJ, Ramani P, Premkumar P, Anuja N. Cytomegalovirus and Mucoepidermoid carcinoma: A possible causal relationship? A pilot study. *J Oral MaxillofacPathol.* 2015 Sep;19(3):319–24.
30. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, NavarasampattiSivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent.* 2019 Jun;28(3):289–95.
31. Ariga P, Nallaswamy D, Jain AR, Ganapathy DM. Determination of correlation of width of Maxillary Anterior Teeth using Extraoral and Intraoral Factors in Indian Population: A systematic review. *World J Dent.* 2018 Feb;9(1):68–75.
32. Kannan A, Venugopalan S. A systematic review on the effect of use of impregnated retraction cords on gingiva. *J Adv Pharm Technol Res.* 2018;11(5):2121.

33. Basha FYS, Ganapathy D, Venugopalan S. Oral hygiene status among pregnant women. *J Adv Pharm Technol Res.* 2018;11(7):3099.
34. Rajakeerthi, Ms N. Natural Product as the Storage medium for an avulsed tooth – A Systematic Review. *CumhurÜnivDişHekimFakderg.* 2019 Jun 11;22(2):249–56.
35. Teja KV, Ramesh S, Priya V. Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study. *J Conserv Dent.* 2018 Nov;21(6):592–6.
36. Menon S, Ks SD, R S, S R, S VK. Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism. *Colloids Surf B Biointerfaces.* 2018 Oct 1;170:280–92.
37. Siddique R, Sureshbabu NM, Somasundaram J, Jacob B, Selvam D. Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi. *J Conserv Dent.* 2019 Jan;22(1):40–7.
38. Nandakumar M, Nasim I. Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis. *J Conserv Dent.* 2018 Sep;21(5):516–20.
39. Manohar MP, Sharma S. A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists. *Indian J Dent Res.* 2018 Nov;29(6):716–20.
40. Hema Shree K, Ramani P, Sherlin H, Sukumaran G, Jeyaraj G, Don KR, et al. Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma - a Systematic Review with Meta Analysis. *PatholOncol Res.* 2019 Apr;25(2):447–53.
41. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative evaluation of remineralizing potential of a paste containing bioactive glass and a topical cream containing casein phosphopeptide-amorphous calcium phosphate: An in vitro study. *Pesqui Bras Odontopediatria Clin Integr.* 2019;19(1):1–10.
42. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol.* 2019 Jun;38(6):694–702.
43. Hussainy SN, Nasim I, Thomas T, Ranjan M. Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up. *J Conserv Dent.* 2018 Sep;21(5):510–5.
44. Hannah R, Ramani P, Herald. J. Sherlin, Ranjith G, Ramasubramanian A, Jayaraj G, et al.

- Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens. *J Adv Pharm Technol Res.* 2018;11(3):1012.
45. Sharma P, Mehta M, Dhanjal DS, Kaur S, Gupta G, Singh H, et al. Emerging trends in the novel drug delivery approaches for the treatment of lung cancer. *Chem Biol Interact.* 2019 Aug 25;309:108720.
 46. Ravinthar K, Jayalakshmi. Recent advancements in laminates and veneers in dentistry. *J Adv Pharm Technol Res.* 2018;11(2):785.
 47. Jose J, Ajitha, Subbaiyan H. Different treatment modalities followed by dental practitioners for Ellis class 2 fracture – A questionnaire-based survey. *Open Dent J.* 2020 Feb 18;14(1):59–65.
 48. Sekar D, Lakshmanan G, Mani P, Biruntha M. Methylation-dependent circulating microRNA 510 in preeclampsia patients. *Hypertens Res.* 2019 Oct;42(10):1647–8.
 49. Kumar D, Antony SDP. Calcified canal and negotiation-A review. *J Adv Pharm Technol Res.* 2018;11(8):3727.
 50. Johnson J, Lakshmanan G, M B, R M V, Kalimuthu K, Sekar D. Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH. *Hypertens Res.* 2020 Apr;43(4):360–2.
 51. Janani K, Palanivelu A, Sandhya R. Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality: an in vivo study. *Braz Dent Sci [Internet].* 2020 Jan 31;23(1). Available from: <https://bds.ict.unesp.br/index.php/cob/article/view/1805>
 52. Seppan P, Muhammed I, Mohanraj KG, Lakshmanan G, Premavathy D, Muthu SJ, et al. Therapeutic potential of *Mucuna pruriens* (Linn.) on ageing induced damage in dorsal nerve of the penis and its implication on erectile function: an experimental study using albino rats. *Aging Male.* 2018 Feb 15;1–14.
 53. Jeevanandan G, Govindaraju L. Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial. *Eur Arch Paediatr Dent.* 2018 Aug;19(4):273–8.
 54. Nandhini JST, Babu KY, Mohanraj KG. Size, shape, prominence and localization of gerdy's tubercle in dry human tibial bones. *J Adv Pharm Technol Res.* 2018;11(8):3604.
 55. Hook CR, Comer RW, Trombly RM, Guinn JW, Shrout MK. Treatment planning processes

- in dental schools. *J Dent Educ.* 2002;66(1):68–74.
56. Berkey DB, Berg RG, Ettinger RL, Mersel A, Mann J. THE OLD-OLD DENTAL PATIENT: THE CHALLENGE OF CLINICAL DECISION-MAKING. *The Journal of the American Dental Association.* 1996 Mar 1;127(3):321–32.
 57. Cohen SR, Orenstein JH. The use of attachments in combination implant and natural-tooth fixed partial dentures: a technical report. *Int J Oral Maxillofac Implants* [Internet]. 1994;9(2). Available from: <http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=08822786&AN=38012782&h=ZsXhBn1%2FVy%2FB%2BFa3VcUHfaRXhiw3WYcB4KgXvG0z%2BWOOkZ9W41prtcRJJbm%2FriA8iG5Hxwg46w%2Ftakqf%2FbDitw%3D%3D&crl=c>
 58. Seligman DA, Pullinger AG, Solberg WK. The Prevalence of Dental Attrition and its Association with Factors of Age, Gender, Occlusion, and TMJ Symptomatology. *J Dent Res.* 1988 Oct 1;67(10):1323–33.
 59. Eshed V, Gopher A, HersHKovitz I. Tooth wear and dental pathology at the advent of agriculture: New evidence from the Levant. *Am J Phys Anthropol.* 2006 Jun;130(2):145–59.
 60. Ahmad I. Anterior dental aesthetics: Dental perspective. *Br Dent J.* 2005 Aug 1;199(3):135–41.
 61. Vallittu PK, Vallittu ASJ, Lassila VP. Dental aesthetics—a survey of attitudes in different groups of patients. *J Dent.* 1996;24(5):335–8.
 62. Teich ST. Risk Assessment-Based Individualized Treatment (RABIT): a comprehensive approach to dental patient recall. *J Dent Educ.* 2013 Apr;77(4):448–57.
 63. Amin M, Nouri R, ElSalhy M, Shah P, Azarpazhooh A. Caries recurrence after treatment under general anaesthesia for early childhood caries: a retrospective cohort study. *Eur Arch Paediatr Dent.* 2015 Aug 1;16(4):325–31.
 64. Kanegane K, Penha SS, Borsatti MA, Rocha RG. Dental anxiety in an emergency dental service. *Rev SaúdePública.* 2003 Dec;37:786–92.
 65. Moy PK, Medina D, Shetty V, Aghaloo TL. Dental implant failure rates and associated risk factors. *Int J Oral Maxillofac Implants* [Internet]. 2005;20(4). Available from: <http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=08822786&AN=36831255&h=VNmPj68p9zAxdj5l0sHdqJT2barTti2Ww8ft1>

2Li9JWLvd%2B2ukgOJNhjQKnpoX%2BFoVEo37QWNtpp6OG9evoOIg%3D%3D&crl=c

66. Ettinger RL. Treatment planning concepts for the ageing patient. *Aust Dent J*. 2015 Mar 11;60:71–85.
67. Armfield JM, Slade GD, Spencer AJ. Dental fear and adult oral health in Australia. *Community Dent Oral Epidemiol*. 2009 Jun;37(3):220–30.
68. Naidu RS, Lalwah S. Dental anxiety in a sample of West Indian adults. *West Indian Med J* [Internet]. 2010;59(5). Available from: https://www.mona.uwi.edu/fms/wimj/system/files/article_pdfs/dr_naidu_wimj.qxd_.pdf
69. Mettes D. Insufficient evidence to support or refute the need for 6-monthly dental check-ups [Internet]. Vol. 6, *Evidence-Based Dentistry*. 2005. p. 62–3. Available from: <http://dx.doi.org/10.1038/sj.ebd.6400341>
70. Collett HA. Influence of dentist-patient relationship on attitudes and adjustment to dental treatment. *The Journal of the American Dental Association*. 1969 Oct 1;79(4):879–84.
71. Patel S, Bay RC, Glick M. A systematic review of dental recall intervals and incidence of dental caries. *J Am Dent Assoc*. 2010 May;141(5):527–39.
72. Jayaraj G, Ramani P, Herald J. Sherlin, Premkumar P, Anuja N. Inter-observer agreement in grading oral epithelial dysplasia – A systematic review [Internet]. Vol. 27, *Journal of Oral and Maxillofacial Surgery, Medicine, and Pathology*. 2015. p. 112–6. Available from: <http://dx.doi.org/10.1016/j.ajoms.2014.01.006>
73. VijayashreePriyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol*. 2019 Dec;90(12):1441–8.
74. Pc J, Marimuthu T, Devadoss P. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res* [Internet]. 2018; Available from: <https://europepmc.org/article/med/29624863>
75. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol*. 2018 Oct;89(10):1241–8.
76. Tefera, B.N., Kim, Y.-D. Ethnobotanical study of Medicinal plants used as Antimalarial and Repellent by Sidama people of Hawassa Zuria district, Southern Ethiopia(2019) *Journal of*

Complementary Medicine Research, 10 (1), pp. 13-26.

77. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygiumcumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. J Oral Pathol Med. 2019 Feb;48(2):115–21.
78. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: Randomized controlled trial. Clin Oral Investig. 2020;1–6.
79. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? Int J Paediatr Dent. 2021 Mar;31(2):285–6.
80. R H, Hannah R, Ramani P, Ramanathan A, R JM, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Vol. 130, Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology. 2020. p. 306–12. Available from: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
81. Chandrasekar R, Chandrasekhar S, Sundari KKS, Ravi P. Development and validation of a formula for objective assessment of cervical vertebral bone age. Prog Orthod. 2020 Oct 12;21(1):38.
82. VijayashreePriyadharsini J, SmilineGirija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen A. baumannii and related species. Arch Oral Biol. 2018 Oct;94:93–8.

Tables:

Table 1: Table depicting the descriptive statistics of percentage of treatment completed and percentage of treatment pending. Majority of the treatments planned (60.05%) were completed followed by (39.95%) pending treatment yet to be completed.

	Percentage of treatment completed	Percentage of treatment pending
Mean	60.05	39.95
Standard Error of Mean	3.496	3.496
Standard Deviation	66.67	33.33

Figure Legends:

Figure 1: Pie chart depicting normal case and emergency case distribution among the study population. Blue represents emergency cases and green represents normal cases. Majority of the cases were normal cases (67%) followed by (33%) emergency cases which required immediate treatment.

Figure 2: Bar graph depicting the percentage of total treatment planned. X axis represents the total number of treatment procedures. Y axis represents the percentage of treatment planned. Blue represents less than 5 treatment procedures, green represents 6-10 treatment procedures, beige represents 10-15 procedures, purple represents 16-20 procedures and yellow represents more than 20 procedures. The association was done using chi square analysis $p=.117$ (statistically not significant). Majority of the study population had less than 5 treatment procedures planned (56%) followed by 6-10 procedures (29%), 10-15 procedures (9%), 16-20 procedures (3%) and more than 20 procedures was only by (3%) of the study population.

Figure 3: Bar graph depicting the association between number of treatment procedures planned with emergency and normal cases. X axis represents the type of cases. Y axis represents the percentage of treatment planned. Blue represents less than 5 treatment procedures, green represents 6-10 treatment procedures, beige represents 10-15 procedures, purple represents 16-20 procedures and yellow represents more than 20 procedures. Though the majority of the normal visit patients had <5 procedures planned (40%), there was no statistically significant difference in the number of planned procedures between normal visit and emergency patients. Pearson chi square, $p=.117$ ($p>0.05$).

Figure 4: Bar graph depicting percentage of total treatment completed. X axis represents the Percentage of treatment completed. Y axis represents the corresponding percentage. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. The association was done using chi square analysis $p=.273$ (statistically not significant). Majority of the study population (26%) had 100 percent treatment completed and (12%) of the study population had 0% treatment completed.

Figure 5: Bar graph depicting the association of percentage of treatment completed with emergency and normal cases. X axis represents the percentage of treatment completed. Y axis represents the count of emergency and normal cases. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. Percentage of treatment completed in

emergency cases (33%) was comparatively lesser than percentage of treatment completed in normal cases (67%). This was however statistically not significant. Pearson Chi square, $p=0.273(p>0.05)$.

Figure 6: Bar graph depicting the association of percentage of treatment completed with total treatment planned. X axis represents the number of treatments planned. Y axis represents the percentage of treatment completed. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. The association was done using chi square analysis 100 % of the planned treatment was completed majorly in cases with less than 5 treatment procedures (20%). However, this was statistically not significant. Pearson's Chi square, $p=0.099(p>0.05)$.




Figure 1: Pie chart depicting normal case and emergency case distribution among the study population. Blue represents emergency cases and green represents normal cases. Majority of the cases were normal cases (67%) followed by (33%) emergency cases which required immediate treatment.




Figure 2: Bar graph depicting the percentage of total treatment planned. X axis represents the total number of treatment procedures. Y axis represents the percentage of treatment planned. Blue represents less than 5 treatment procedures, green represents 6-10 treatment procedures, beige represents 10-15 procedures, purple represents 16-20 procedures and yellow represents more than 20 procedures. The association was done using chi square analysis $p=.117$ (statistically not significant). Majority of the study population had less than 5 treatment procedures planned (56%) followed by 6-10 procedures (29%), 10-15 procedures (9%), 16-20 procedures (3%) and more than 20 procedures was only by (3%) of the study population.




Figure 3: Bar graph depicting the association between number of treatment procedures planned with emergency and normal cases. X axis represents the type of cases. Y axis represents the percentage of treatment planned. Blue represents less than 5 treatment procedures, green represents 6-10 treatment procedures, beige represents 10-15 procedures, purple represents 16-20 procedures and yellow represents more than 20 procedures. Though the majority of the normal visit patients had <5 procedures planned (40%), there was no statistically significant difference in the number of planned procedures between normal visit and emergency patients. Pearson chi square, $p=.117$ ($p>0.05$).





Figure 4:Bar graph depicting percentage of total treatment completed.X axis represents the Percentage of treatment completed.Y axis represents the corresponding percentage. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. The association was done using chi square analysis $p=.273$ (statistically not significant).majority of the study population (26%) had 100 percent treatment completed and (12%)of the study population had 0% treatment completed.

Figure 5: Bar graph depicting the association of percentage of treatment completed with emergency and normal cases. X axis represents the percentage of treatment completed . Y axis represents the count of emergency and normal cases. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. Percentage of treatment completed in emergency cases (33%) was comparatively lesser than percentage of treatment completed in normal cases (67%). This was however statistically not significant. Pearson Chi square , $p=0.273(p>0.05)$.






Figure 6: Bar graph depicting the association of percentage of treatment completed with total treatment planned. X axis represents the number of treatments planned . Y axis represents the percentage of treatment completed. Blue represents less than 25% treatment completed, green represents 26-50% treatment completed, beige represents 51-75% treatment completed, purple represents 76-99% treatment completed, yellow represents 100% treatment completed and red represents 0% treatment completed. The association was done using chi square analysis 100 % of the planned treatment was completed majorly in cases with less than 5 treatment procedures(20%) . However, this was statistically not significant. Pearson's Chi square, $p=0.099(p>0.05)$.