

Prevalence of Oral Mucosal Lesions among Tobacco Users Visiting a Dental Hospital - A Retrospective Study

Authors: Oviya. V. J¹, Pradeep Kumar. R², Deepa Gurunathan³

Type of manuscript: Retrospective study

Running title : Prevalence of oral mucosal lesions among tobacco users visiting a dental hospital

V. J. Oviya,

Saveetha Dental College,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
162, Poonamallee High Road, Chennai - 600077, Tamil Nadu, India
Email id : 151501075.sdc@saveetha.com

Pradeep kumar.R

Professor and Head,
Department of Public Health Dentistry,
Saveetha Dental College,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
162, Poonamallee High Road, Chennai - 600077, Tamil Nadu, India
Email id : pradeepkumarr.sdc@saveetha.com

Deepa Gurunathan,

Professor and Head,
Department of Pedodontics and preventive dentistry,
Saveetha Dental College,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
162, Poonamallee High Road, Chennai - 600077, Tamil Nadu, India
Email id : deepag@saveetha.com

Corresponding author:

Pradeep kumar.R

Professor and Head,
Department of Public Health Dentistry,
Saveetha Dental College,
Saveetha Institute of Medical and Technical Sciences,
Saveetha University,
162, Poonamallee High Road, Chennai - 600077, Tamil Nadu, India

Email id :pradeepkumarr.sdc@saveetha.com

Ph no.: 99403 91698

ABSTRACT:

Oral Mucosal lesions(OMLs) are any alteration in oral mucosal surfaces which can cause discomfort and pain. The aim of this study is to analyse the common oral mucosal lesions among tobacco users visiting dental hospital. A retrospective study was conducted by evaluating and analysing 1113 patient case records visiting a dental hospital from June(2019) to March(2020) who were tobacco users. Data such as age, gender, type of tobacco, site and type of lesions were documented. Descriptive analysis and Chi Square test were done. The result showed that prevalence of oral lesions were higher among patients of age group 20-40 years(50.6%) and among males patients(96.6%). Leukoplakia(25.6%) and tobacco pouch keratosis(25.6%) were commonly reported lesions. Among the patients who had presented with oral lesions, 87.3% of the tobacco users had only single oral lesions and buccal mucosa(63.7%) being the most common site of occurrence. The present study provides epidemiological information on OMLs among patients seeking dental care, which could be a valuable source for future community tobacco control programs .

Keywords:

Oral mucosal lesions, Prevalence, Potential malignant disorders, Smoking, Tobacco, Patients, Smokeless

INTRODUCTION:

Smokeless tobacco use is more common in India and SouthEast Asia(Organization and Others, 2016). India is the second largest producer and consumer of tobacco next to China(Kaur and Jain, 2011). Tobacco contains numerous carcinogens mostly polycyclic aromatic hydrocarbons, aromatic amines and nitrosamines which cause cellular damages to the host body(Mishra *et al.*, 2014). Tobacco usage is one of the main risk factors for causing cardiovascular diseases and lung diseases(Harini and Leelavathi, 2019)

In India, the use of cigarette smoking is comparatively lesser than the usage of smokeless tobacco as it is more socially acceptable and can be practiced without detection(Gupta *et al.*, 1980). The WHO also predicts that India will have deaths attributed to tobacco usage in future years and highest rates of oral cancer due to the prevalence of tobacco chewing(World Health Organization, 1997).

Use of tobacco in the form of either smokeless or smoking has been highly associated with oral mucosal lesions, which in turn may eventually become malignant(Reddy *et al.*, 2015). Oral mucosal lesions are due to any abnormal alteration in colour, surfaces, aspect, swelling and integrity of oral mucosa. Though many are benign lesions, some may cause significant pathology

and malignancy. These lesions can affect daily life functions through impairment in mastication, speech and swallowing with symptoms of pain, burning and irritants(Feng *et al.*, 2015).

In comparison to western population in oral cancer represents about 3% of malignancies, 30% of all cancers occurred in world accounts to India. This is due to regional variation in the prevalence and pattern of habits(Dangi, Kinnunen and Zavras, 2012). The treatment needs also vary with every geographic region within a country and even between different communities.

Previously our team has conducted numerous cross sectional studies(Prabakar, John and Srisakthi, 2016; Kannan *et al.*, 2017; Neralla *et al.*, 2019), clinical trials(Prabakar, John, Arumugham, Kumar and Sakthi, 2018b; Prabakar, John, Arumugham, Kumar and Srisakthi, 2018; Khatri *et al.*, 2019; Pratha and Prabakar, 2019; Mathew *et al.*, 2020a; Samuel, Acharya and Rao, 2020), in-vitro studies(Kumar and Preethi, 2017; Kumar and Vijayalakshmi, 2017; Prabakar, John, Arumugham, Kumar and Sakthi, 2018a; Mohapatra *et al.*, 2019; Pavithra and Jayashri, 2019), over the past 5 years. Now we are focussing on epidemiological surveys. Our team has rich experience in research and we have collaborated with numerous authors over various topics in the past decade (Ariga *et al.*, 2018; Basha, Ganapathy and Venugopalan, 2018; Hannah *et al.*, 2018; Hussainy *et al.*, 2018; Jeevanandan and Govindaraju, 2018; Kannan and Venugopalan, 2018; Kumar and Antony, 2018; Manohar and Sharma, 2018; Menon *et al.*, 2018; Nandakumar and Nasim, 2018; Nandhini, Babu and Mohanraj, 2018; Ravinthar and Jayalakshmi, 2018; Seppan *et al.*, 2018; Teja, Ramesh and Priya, 2018; Duraisamy *et al.*, 2019; Gheena and Ezhilarasan, 2019; Hema Shree *et al.*, 2019; Rajakeerthi and Ms, 2019; Rajendran *et al.*, 2019; Sekar *et al.*, 2019; Sharma *et al.*, 2019; Siddique *et al.*, 2019; Janani, Palanivelu and Sandhya, 2020; Johnson *et al.*, 2020; Jose, Ajitha and Subbaiyan, 2020).

Since the epidemiological studies provide information which are essential to understand prevalence and severity of oral diseases in specific populations, the present study was carried out to study and analyse prevalence of oral mucosal lesions among tobacco users visiting a dental hospital in Chennai.

MATERIALS AND METHODS:

Sampling:

This study was conducted in a university setting. The study samples were chosen from the patients visiting a hospital in Chennai from June(2019) to March(2020).

Data collection:

The retrospective study was carried out among patients of all age groups. Data collection was done through reviewing the records of 86000 patients between June(2019) - March(2020). A total of 1113 patients reported with tobacco habits for at least a year with complete data. Data

such as the patient's age, gender, type of tobacco, site and type of lesions were noted. The data collected were cross verified with intraoral photographs.

Inclusion criteria:

Only the patients who were using tobacco for at least a year of all age groups were included.

Exclusion criteria:

Patients with severe systemic diseases and under special care were excluded. Incomplete/ censored data were excluded too.

Approval:

Ethical clearance was obtained from the Institutional Scientific Review Board of the SaveethaUniversity(SDC/SIHEC/2020/DIASDATA/0619-0320).

Data analysis:

The data collected was entered in the excel sheet and transferred to SPSS software. Data was analysed using SPSS software though frequency distribution and Chi-square tests.

RESULTS:

A total of 1113 reported to the hospital with tobacco habits. The mean age of the study population was 38.1years. Among them, 1084(97.3%) of the patients were males and 29(2.5%) were females.

In the present study, usage of smoking was prevalent among all age groups. Smokeless tobacco use was the most common habit among females(58.7%) and smoking was the most common habit among males(76.8%)(Table 1).

Majority of the study group belong to the age of 20-40(58.4%) with tobacco usage which is however not statistically significant(Figure 1). In this study, 75.9% of the study population were smokers, 16.0% were smokeless tobacco users, 8% used both smoking and smokeless forms of tobacco.

In the present study, 20.5% of the total patients presented with single or multiple oral mucosal lesions. The prevalence of oromucosallesions were more among males(96.6%) than females(3.4%). About 27.5% of females and 22.3% of the males reported with one or more oromucosal lesions. (Figure 2).

In the present study, oromucosal lesions were more common among 20-40 years old (50.6%)(Figure 3). Out of 20.5% of the patients who had reported with oral lesions, it was found that about 87.3% of the patients had only one mucosal lesion, 9.3% of the patients had two lesions and only 3.4% of the patients reported with more than two lesions.

Leukoplakia(25.6%), Tobacco pouch keratosis (25.6%) and OSMF (13.9%) were the most common lesions diagnosed in this study(Figure 4). Tobacco pouch keratosis (15.8%) was the most common in the age group of 20-40 years old while leukoplakia was more prevalent among patients of all other age groups(Figure 5). Among them, higher prevalence was seen at the site of buccal mucosa (63.7%) followed by labial mucosa (10.4%) and hard palate (10.1%)(Figure 6).

DISCUSSION:

Oromucosal lesions and diseases are caused by infections, systemic diseases, drugs, adverse factors like tobacco, betel nut or alcohol consumption(Pindborg, 1968). Tobacco usage is also positively associated which have the potential for malignant transformation(Saraswathi *et al.*, 2006).

The overall prevalence of oromucosal lesions among tobacco users in this study was 20.5%. The prevalence ratio is much higher than the studies conducted by Saraswathi *et al.*(Saraswathi *et al.*, 2006), Chung *et al.*(Chung *et al.*, 2005) and Mani *et al.*(Mani, 1985). However few studies have reported higher prevalence rates by Bhowate *et al.*(Bhowate *et al.*, 1994) and Patil *et al.*(Patil, Bathi and Chaudhari, 2013). The difference in this finding may be due to the different methodologies used by different researchers. Besides, due to variations in cultural and social impacts, it is difficult to compare these studies.

In India, adult males are expected to smoke, while smoking in females is considered a taboo. This is greatly mirrored in the results, where 97.3% of the participants were male. This is in accordance with many studies(Lin, Corbet and Lo, 2001; Rani *et al.*, 2003; Pentenero, Broccoletti and Carbone, 2008; Vellappally *et al.*, 2008). In the present study, only 1.2% of the smokers were females, 98.7% were males. About 48.3% of females have the habit of chewing tobacco. The results are similar to the findings reported by Vellappally *et al.*(Vellappally *et al.*, 2008) and Rani *et al.*(Rani *et al.*, 2003). Though smoking among young people and females is considered taboo in India, there is no such restriction against the usage of smokeless tobacco as it is easy to practice without any detection,

In the current study, tobacco usage was highly prevalent among young adults of age group 20-40 years compared to others. This is in accordance with studies done by Sunil *et al.*(Mishra *et al.*, 2014), Vellappally *et al.*(Vellappally *et al.*, 2008)), Abhishek *et al.*(Abhishek *et al.*,

2012). The presence of lesions was more prevalent among the age group of 20-40 years showing strong association with tobacco usage. About 58.6% of the smokeless tobacco users belong to this 20-40 years of age group. This finding is similar to the studies done by Singh et al (Singh and Ladusingh, 2014) and Mohan et al (Mohan, Lando and Panneer, 2018) who reported that smokeless tobacco usage was higher in younger adults.

In this study, 27.5% of the females and 22.3% of the males in the sample presented with lesions. However, oromucosal lesions were more prevalent among males (96.6%) compared to females (3.3%). Previous studies by Lin et al (Lin, Corbet and Lo, 2001), Pentenero et al (Pentenero, Broccolotti and Carbone, 2008) and Patil et al (Patil, Doni and Maheshwari, 2015) reported similar findings. However few studies reported female predominance (Majeed and Abid, 2009; Al-Gburi and Mudhir, 2018). This is due to the cultural impacts in different populations.

In the present study, Leukoplakia (25.6%) and tobacco pouch keratosis (25.6%) were the most commonly reported lesions followed by oral submucous fibrosis (13.9%). Many studies reported leukoplakia as the most commonest oral mucosal lesion (Ikeda *et al.*, 1991; Sujatha, Hebbar and Pai, 2012; Al-Attas *et al.*, 2014). In contrast few studies reported less prevalence of leukoplakia (Bansal, Sogi and Veerasha, 2010).

Many studies have reported oral submucous fibrosis as the commonest lesion among smokers (Patil, Bathi and Chaudhari, 2013; Reddy *et al.*, 2015; Aishwarya *et al.*, 2017; Alshayeb, Mathew and Varma, 2019). Few studies reported higher prevalence of tobacco pouch keratosis (Joshi and Tailor, 2016; Kamala *et al.*, 2019).

In the present study, the tobacco pouch keratosis (15.8%) was common among patients of 20-40 years age group, while leukoplakia was commonly presented in all other age groups. This could be due to the increased use of chewing tobacco among young adults. This is in accordance with similar studies done by Singh et al and Mohan et al (Singh and Ladusingh, 2014; Mohan, Lando and Panneer, 2018).

The oral mucosal lesions were more prevalent at buccal mucosa (63.7%). This finding is supported by few other studies (Ahmadi-Motamaye *et al.*, 2013; Patil, Bathi and Chaudhari, 2013; Krishna Priya, Srinivas and Devaki, 2018). Our institution is passionate about high quality evidence based research and has excelled in various fields (Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; Vijayashree Priyadharsini, Smiline Girija and Paramasivam, 2018; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Ramadurai *et al.*, 2019; Sridharan *et al.*, 2019; Vijayashree Priyadharsini, 2019; Chandrasekar *et al.*, 2020; Mathew *et al.*, 2020b; R *et al.*, 2020; Samuel, 2021).

The present study had few limitations like histopathological analysis not included. But still the present was able to provide the prevalence of oral mucosal lesions among tobacco users visiting a dental hospital.

CONCLUSION:

Within the limits of the study, the prevalence of oromucosal lesions was higher among tobacco users. Many of the tobacco related oromucosal lesions are asymptomatic and go unnoticed till they reach their higher stages. The lack of awareness about the harmful effects of tobacco and the ignorance towards its effects was the major cause for failure towards early diagnosis and prompt treatment. The present study provides epidemiological information on oral mucosal lesions among patients seeking dental care, which could be a valuable source for future community tobacco control programs.

AUTHOR'S CONTRIBUTION: All authors contributed to the design and implementation of the research, analysis of the results and to the writing of the manuscript.

ACKNOWLEDGEMENT- Nil

CONFLICT OF INTEREST:

The authors declared that they have no conflicts of interest.

REFERENCES:

- [1]. Abhishek, K. *et al.* (2012) 'Oral premalignant lesions associated with areca nut and tobacco chewing among the tobacco industry workers in area of rural Maharashtra', *National J Community Med*, 3, pp. 333–338.
- [2]. Ahmadi-Motamayel, F. *et al.* (2013) 'Prevalence of oral mucosal lesions in male smokers and nonsmokers', *Chonnam medical journal*, 49(2), pp. 65–68.
- [3]. Aishwarya, K. M. *et al.* (2017) 'Effect of Frequency and Duration of Tobacco Use on Oral Mucosal Lesions – A Cross-Sectional Study among Tobacco Users in Hyderabad, India', *Asian Pacific journal of cancer prevention: APJCP*, 18(8), pp. 2233–2238.
- [4]. Al-Attas, S. A. *et al.* (2014) 'Prevalence of potentially malignant oral mucosal lesions among tobacco users in Jeddah, Saudi Arabia', *Asian Pacific journal of cancer prevention: APJCP*, 15(2), pp. 757–762.
- [5]. Al-Gburi, S. M. and Mudhir, S. H. (2018) 'The Prevalence of the Oral Mucosal Lesions among Adult Patients in Abu Ghraib City (Iraq)', *Journal of Research in Medical and Dental Science*, 6(5), pp. 145–148.
- [6]. Alshayeb, M., Mathew, A. and Varma, S. (2019) 'Prevalence and distribution of oral mucosal lesions associated with tobacco use in patients visiting a dental school in Ajman', *Onkologia i.* Available at: <https://www.oncologyradiotherapy.com/abstract/prevalence-and-distribution-of-oral-mucosal-lesions-associated-with-tobacco-use-in-patients-visiting-a-dental-school-in--43916.html>.
- [7]. Ariga, P. *et al.* (2018) 'Determination of correlation of width of Maxillary Anterior Teeth using

- Extraoral and Intraoral Factors in Indian Population: A systematic review', *World journal of dentistry*, 9(1), pp. 68–75.
- [8]. Bansal, V., Sogi, G. M. and Veerasha, K. L. (2010) 'Assessment of oral health status and treatment needs of elders associated with elders' homes of Ambala division, Haryana, India', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 21(2), pp. 244–247.
- [9]. Basha, F. Y. S., Ganapathy, D. and Venugopalan, S. (2018) 'Oral hygiene status among pregnant women', *Journal of advanced pharmaceutical technology & research*, 11(7), p. 3099.
- [10]. Kwok, M., Tran, T.M.L. For the future and possible ensuing waves of COVID-19: A perspective to consider when disseminating data(2020) *Journal of Population Therapeutics and Clinical Pharmacology*, 27 (SP1), pp. e53-e57.
- [11]. Chandrasekar, R. et al. (2020) 'Development and validation of a formula for objective assessment of cervical vertebral bone age', *Progress in orthodontics*, 21(1), p. 38.
- [12]. Chung, C.-H. et al. (2005) 'Oral precancerous disorders associated with areca quid chewing, smoking, and alcohol drinking in southern Taiwan', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 34(8), pp. 460–466.
- [13]. Dangi, J., Kinnunen, T. H. and Zavras, A. I. (2012) 'Challenges in global improvement of oral cancer outcomes: findings from rural Northern India', *Tobacco induced diseases*, 10, p. 5.
- [14]. Duraisamy, R. et al. (2019) 'Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments', *Implant dentistry*, 28(3), pp. 289–295.
- [15]. Ezhilarasan, D., Apoorva, V. S. and Ashok Vardhan, N. (2019) 'Syzygiumcumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121.
- [16]. Feng, J. et al. (2015) 'Prevalence and distribution of oral mucosal lesions: a cross-sectional study in Shanghai, China', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 44(7), pp. 490–494.
- [17]. Gheena, S. and Ezhilarasan, D. (2019) 'Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells', *Human & experimental toxicology*, 38(6), pp. 694–702.
- [18]. Gupta, P. C. et al. (1980) 'Incidence rates of oral cancer and natural history of oral precancerous lesions in a 10-year follow-up study of Indian villagers', *Community dentistry and oral epidemiology*, 8(6), pp. 283–333.
- [19]. Hannah, R. et al. (2018) 'Awareness about the use, ethics and scope of dental photography among undergraduate dental students dentist behind the lens', *Journal of advanced pharmaceutical technology & research*, 11(3), p. 1012.
- [20]. Harini, G. and Leelavathi, L. (2019) 'Nicotine Replacement Therapy for Smoking Cessation-An Overview', *Indian Journal of Public Health Research & Development*, 10(11), pp. 3588–3592.

- [21]. Hema Shree, K. *et al.* (2019) 'Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma - a Systematic Review with Meta Analysis', *Pathology oncology research: POR*, 25(2), pp. 447–453.
- [22]. Hussainy, S. N. *et al.* (2018) 'Clinical performance of resin-modified glass ionomer cement, flowable composite, and polyacid-modified resin composite in noncarious cervical lesions: One-year follow-up', *Journal of conservative dentistry: JCD*, 21(5), pp. 510–515.
- [23]. Ikeda, N. *et al.* (1991) 'Epidemiological study of oral leukoplakia based on mass screening for oral mucosal diseases in a selected Japanese population', *Community dentistry and oral epidemiology*, 19(3), pp. 160–163.
- [24]. Janani, K., Palanivelu, A. and Sandhya, R. (2020) 'Diagnostic accuracy of dental pulse oximeter with customized sensor holder, thermal test and electric pulp test for the evaluation of pulp vitality: an in vivo study', *Brazilian dental science*, 23(1). doi: 10.14295/bds.2020.v23i1.1805.
- [25]. Jeevanandan, G. and Govindaraju, L. (2018) 'Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial', *European archives of paediatric dentistry: official journal of the European Academy of Paediatric Dentistry*, 19(4), pp. 273–278.
- [26]. Johnson, J. *et al.* (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH', *Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362.
- [27]. Jose, J., Ajitha and Subbaiyan, H. (2020) 'Different treatment modalities followed by dental practitioners for Ellis class 2 fracture – A questionnaire-based survey', *The open dentistry journal*, 14(1), pp. 59–65.
- [28]. Joshi, M. and Tailor, M. (2016) 'Prevalence of most commonly reported tobacco-associated lesions in central Gujarat: A hospital-based cross-sectional study', *Indian journal of dental research: official publication of Indian Society for Dental Research*. Available at: <http://www.ijdr.in/article.asp?issn=0970-9290;year=2016;volume=27;issue=4;spage=405;epage=409;aulast=Joshi>.
- [29]. Kamala, K. A. *et al.* (2019) 'Prevalence of oromucosal lesions in relation to tobacco habit among a Western Maharashtra population', *Indian journal of cancer*, 56(1), pp. 15–18.
- [30]. Kannan, A. and Venugopalan, S. (2018) 'A systematic review on the effect of use of impregnated retraction cords on gingiva', *Journal of advanced pharmaceutical technology & research*, 11(5), p. 2121.
- [31]. Kannan, S. S. D. *et al.* (2017) *AWARENESS AND ATTITUDE TOWARDS MASS DISASTER AND ITS MANAGEMENT AMONG HOUSE SURGEONS IN A DENTAL COLLEGE AND HOSPITAL IN CHENNAI, INDIA*. WIT Press, pp. 121–129.
- [32]. Kaur, J. and Jain, D. C. (2011) 'Tobacco control policies in India: implementation and challenges', *Indian journal of public health*, 55(3), pp. 220–227.
- [33]. Khatri, S. G. *et al.* (2019) 'Retention of moisture-tolerant fluoride-releasing sealant and amorphous calcium phosphate-containing sealant in 6-9-year-old children: A randomized controlled trial', *Journal of the Indian Society of Pedodontics and Preventive Dentistry*, 37(1),

pp. 92–98.

- [34]. Krishna Priya, M., Srinivas, P. and Devaki, T. (2018) 'Evaluation of the Prevalence of Oral Mucosal Lesions in a Population of Eastern Coast of South India', *Journal of International Society of Preventive & Community Dentistry*, 8(5), pp. 396–401.
- [35]. Kumar, D. and Antony, S. D. P. (2018) 'Calcified canal and negotiation-A review', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3727.
- [36]. Kumar, R. P. and Preethi, R. (2017) 'Assessment of Water Quality and Pollution of Porur, Chembarambakkam and Puzhal Lake', *Journal of pharmacy research*. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=10&issue=7&article=032>.
- [37]. Kumar, R. P. and Vijayalakshmi, B. (2017) 'Assessment of fluoride concentration in ground water in Madurai district, Tamil Nadu, India', *Journal of pharmacy research*. Available at: <http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=10&issue=1&article=063>.
- [38]. Lin, H. C., Corbet, E. F. and Lo, E. C. M. (2001) 'Oral mucosal lesions in adult Chinese', *Journal of dental research*. Available at: <https://journals.sagepub.com/doi/abs/10.1177/00220345010800052001>.
- [39]. Majeed, A. H. and Abid, K. J. (2009) 'Prevalence of oral mucosal lesions in Missan governorate', *Journal of baghdad college of dentistry*. Available at: <https://www.iasj.net/iasj?func=fulltext&aId=1659>.
- [40]. Mani, N. J. (1985) 'Preliminary report on prevalence of oral cancer and precancerous lesions among dental patients in Saudi Arabia', *Community dentistry and oral epidemiology*, 13(4), pp. 247–248.
- [41]. Manohar, M. P. and Sharma, S. (2018) 'A survey of the knowledge, attitude, and awareness about the principal choice of intracanal medicaments among the general dental practitioners and nonendodontic specialists', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(6), pp. 716–720.
- [42]. Mathew, M. G. *et al.* (2020a) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial', *Clinical oral investigations*. doi: 10.1007/s00784-020-03204-9.
- [43]. Mathew, M. G. *et al.* (2020b) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: Randomized controlled trial', *Clinical oral investigations*, pp. 1–6.
- [44]. Menon, S. *et al.* (2018) 'Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism', *Colloids and surfaces. B, Biointerfaces*, 170, pp. 280–292.
- [45]. Mishra, S. S. *et al.* (2014) 'Prevalence of oral premalignant lesions and conditions in patients with tobacco and tobacco-related habits reporting to a dental institution in Aurangabad', *Journal of Indian Academy of Oral Medicine and Radiology*, 26(2), p. 152.
- [46]. Mohan, P., Lando, H. A. and Panneer, S. (2018) 'Assessment of tobacco consumption and control in India', *Indian Journal of Clinical*. Available at: <https://journals.sagepub.com/doi/abs/10.1177/1179916118759289>.

- [47]. Mohapatra, S. *et al.* (2019) 'Assessment of Microhardness of Enamel Carious Like Lesions After Treatment with Nova Min, Bio Min and Remin Pro Containing Toothpastes: An in Vitro Study', *Indian Journal of Public Health Research & Development*, 10(10), pp. 375–380.
- [48]. Nandakumar, M. and Nasim, I. (2018) 'Comparative evaluation of grape seed and cranberry extracts in preventing enamel erosion: An optical emission spectrometric analysis', *Journal of conservative dentistry: JCD*, 21(5), pp. 516–520.
- [49]. Nandhini, J. S. T., Babu, K. Y. and Mohanraj, K. G. (2018) 'Size, shape, prominence and localization of gerdy's tubercle in dry human tibial bones', *Journal of advanced pharmaceutical technology & research*, 11(8), p. 3604.
- [50]. Neralla, M. *et al.* (2019) 'Role of nutrition in rehabilitation of patients following surgery for oral squamous cell carcinoma', *International Journal of Research in Pharmaceutical Sciences*, 10(4), pp. 3197–3203.
- [51]. Organization, W. H. and Others (2016) '90% of smokeless tobacco users live in South-East Asia'.
- [52]. Patil, P. B., Bathi, R. and Chaudhari, S. (2013) 'Prevalence of oral mucosal lesions in dental patients with tobacco smoking, chewing, and mixed habits: A cross-sectional study in South India', *Journal of family & community medicine*, 20(2), pp. 130–135.
- [53]. Patil, S., Doni, B. and Maheshwari, S. (2015) 'Prevalence and distribution of oral mucosal lesions in a geriatric Indian population', *Canadian geriatrics journal: CGJ*, 18(1), pp. 11–14.
- [54]. Pavithra, R. P. and Jayashri, P. (2019) 'Influence of Naturally Occurring Phytochemicals on Oral Health', *Research Journal of Pharmacy and Technology*, 12(8), pp. 3979–3983.
- [55]. Pc, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research*. Available at: <https://europepmc.org/article/med/29624863>.
- [56]. Pentenero, M., Broccoletti, R. and Carbone, M. (2008) 'The prevalence of oral mucosal lesions in adults from the Turin area', *Oral-prophylaxe / Herausgeber, Verein fur Zahnhygiene e.V.* Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1601-0825.2007.01391.x>.
- [57]. Pindborg, J. J. (1968) *Atlas of diseases of the oral mucosa*. Munksgaard.
- [58]. Prabakar, J., John, J., Arumugham, I. M., Kumar, R. P. and Srisakthi, D. (2018) 'Comparative Evaluation of Retention, Cariostatic Effect and Discoloration of Conventional and Hydrophilic Sealants - A Single Blinded Randomized Split Mouth Clinical Trial', *Contemporary clinical dentistry*, 9(Suppl 2), pp. S233–S239.
- [59]. Prabakar, J., John, J., Arumugham, I. M., Kumar, R. P. and Sakthi, D. S. (2018a) 'Comparative Evaluation of the Viscosity and Length of Resin Tags of Conventional and Hydrophilic Pit and Fissure Sealants on Permanent Molars: An In vitro Study', *Contemporary clinical dentistry*, 9(3), pp. 388–394.
- [60]. Prabakar, J., John, J., Arumugham, I. M., Kumar, R. P. and Sakthi, D. S. (2018b) 'Comparing the Effectiveness of Probiotic, Green Tea, and Chlorhexidine- and Fluoride-containing Dentifrices on Oral Microbial Flora: A Double-blind, Randomized Clinical Trial', *Contemporary clinical dentistry*, 9(4), pp. 560–569.

- [61]. Prabakar, J., John, J. and Srisakthi, D. (2016) 'Prevalence of dental caries and treatment needs among school going children of Chandigarh', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 27(5), pp. 547–552.
- [62]. Pratha, A. A. and Prabakar, J. (2019) 'Comparing the effect of Carbonated and energy drinks on salivary pH-In Vivo Randomized Controlled Trial', *Journal of pharmacy research*. Available at:
<http://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=10&article=019>.
- [63]. Rajakeerthi and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Üniversitesi Diş Hekimliği Fakültesi Dergisi*, 22(2), pp. 249–256.
- [64]. Rajendran, R. *et al.* (2019) 'Comparative evaluation of remineralizing potential of a paste containing bioactive glass and a topical cream containing casein phosphopeptide-amorphous calcium phosphate: An in vitro study', *Pesquisabrasileira em Odontopediatria e Clínica Integrada*, 19(1), pp. 1–10.
- [65]. Ramadurai, N. *et al.* (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial', *Clinical oral investigations*, 23(9), pp. 3543–3550.
- [66]. Ramesh, A. *et al.* (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248.
- [67]. Rani, M. *et al.* (2003) 'Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross sectional household survey', *Tobacco control*, 12(4), p. e4.
- [68]. Ravinthar, K. and Jayalakshmi (2018) 'Recent advancements in laminates and veneers in dentistry', *Journal of advanced pharmaceutical technology & research*, 11(2), p. 785.
- [69]. Reddy, S. S. *et al.* (2015) 'Prevalence of oral mucosal lesions among chewing tobacco users: A cross-sectional study', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 26(5), pp. 537–541.
- [70]. R, H. *et al.* (2020) 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, pp. 306–312. doi: 10.1016/j.oooo.2020.06.021.
- [71]. Samuel, S. R. (2021) 'Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life?', *International journal of paediatric dentistry / the British Paedodontic Society [and] the International Association of Dentistry for Children*, 31(2), pp. 285–286.
- [72]. Samuel, S. R., Acharya, S. and Rao, J. C. (2020) 'School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial', *Journal of public health dentistry*, 80(1), pp. 51–60.
- [73]. Saraswathi, T. R. *et al.* (2006) 'Prevalence of oral lesions in relation to habits: Cross-sectional study in South India', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 17(3), pp. 121–125.
- [74]. Sekar, D. *et al.* (2019) 'Methylation-dependent circulating microRNA 510 in preeclampsia patients', *Hypertension research: official journal of the Japanese Society of Hypertension*,

42(10), pp. 1647–1648.

- [75]. Seppan, P. *et al.* (2018) ‘Therapeutic potential of *Mucuna pruriens* (Linn.) on ageing induced damage in dorsal nerve of the penis and its implication on erectile function: an experimental study using albino rats’, *The aging male: the official journal of the International Society for the Study of the Aging Male*, pp. 1–14.
- [76]. Sharma, P. *et al.* (2019) ‘Emerging trends in the novel drug delivery approaches for the treatment of lung cancer’, *Chemico-biological interactions*, 309, p. 108720.
- [77]. Siddique, R. *et al.* (2019) ‘Qualitative and quantitative analysis of precipitate formation following interaction of chlorhexidine with sodium hypochlorite, neem, and tulsi’, *Journal of conservative dentistry: JCD*, 22(1), pp. 40–47.
- [78]. Singh, A. and Ladusingh, L. (2014) ‘Prevalence and determinants of tobacco use in India: evidence from recent Global Adult Tobacco Survey data’, *PloS one*, 9(12), p. e114073.
- [79]. Sridharan, G. *et al.* (2019) ‘Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma’, *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306.
- [80]. Sujatha, D., Hebbar, P. B. and Pai, A. (2012) ‘Prevalence and correlation of oral lesions among tobacco smokers, tobacco chewers, areca nut and alcohol users’, *Asian Pacific journal of cancer prevention: APJCP*, 13(4), pp. 1633–1637.
- [81]. Teja, K. V., Ramesh, S. and Priya, V. (2018) ‘Regulation of matrix metalloproteinase-3 gene expression in inflammation: A molecular study’, *Journal of conservative dentistry: JCD*, 21(6), pp. 592–596.
- [82]. Vellappally, S. *et al.* (2008) ‘Tobacco habits and oral health status in selected Indian population’, *Central European journal of public health*, 16(2), pp. 77–84.
- [83]. VijayashreePriyadharsini, J. (2019) ‘In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens’, *Journal of periodontology*, 90(12), pp. 1441–1448.
- [84]. VijayashreePriyadharsini, J., SmilineGirija, A. S. and Paramasivam, A. (2018) ‘In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species’, *Archives of oral biology*, 94, pp. 93–98.
- [85]. World Health Organization (1997) *Tobacco or health : a global status report*. World Health Organization.

GRAPHS AND TABLES:

Gender	Form of tobacco			Total
	Smoking (N %)	Smokeless (N %)	Both (N %)	
Male	833 (76.8%)	162 (15.0)	89 (8.2%)	1084
Female	12 (41.3%)	17 (58.7%)	0 (0%)	29
Total	845 (75.9%)	179 (16.1%)	89 (8.0%)	1113

Table 1: Gender wise distribution of study population based on the form of tobacco usage. Prevalence of smoking was found to be higher among males(76.8%) and smokeless tobacco usage found to be higher among females(58.7%).




Figure 1: Bar chart showing the distribution of study population with tobacco habits based on age group. X-axis shows the age group distribution and Y-axis shows the distribution of the study population based on the form of tobacco usage in percentage. Prevalence of smoking was found to be higher among patients of age group 20-40(43.8%), however it was not statistically significant.(Chi square test; $\chi^2=3.946$, $df=6$, $pValue=0.684(>0.05)$)




Figure 2: Bar chart showing the distribution of study population with the presence of oral lesions according to gender. X-axis shows the gender wise distribution of the study population and Y-axis shows the distribution of the study population with the presence of oral lesions in percentage. Prevalence of oral lesions was found to be higher among males(96.6%-violet) compared to females(3.4%-white).




Figure 3: Bar chart showing the distribution of study population with the presence of oral lesions according to age group. X-axis shows the age wise distribution of the study population and Y-axis shows the distribution of the study population with the presence of oral lesions in percentage. Prevalence of oral lesions was found to be higher among the patients of age group 20-40(50.6%-beige).




Figure 4 : Bar chart showing the distribution of study population with various oral lesions. X-axis shows the distribution of the oral mucosal lesions and Y-axis shows the distribution of the study population presented with oral lesions in percentage. Leukoplakia(25.6%-blue) and tobacco pouch keratosis(25.6%-red) were found to be the most common oral lesions among tobacco users in this study.




Figure 5: Bar chart showing the distribution of study population with various oral mucosal lesions according to age group. X-axis shows age wise distribution of the study population and Y-axis shows the distribution of the study population presented with various oral lesions in percentage. Tobacco pouch keratosis was the most prevalent oral lesion among the patients of the age group 20-40 years(15.8%-red) and leukoplakia was the most common lesion prevalent among the patients of the age group 40-60 years(11.3%-blue). This finding was statistically significant.(Chi square test; $\chi^2=88.514$, $df=60$, $pValue=0.010(<0.05)$)




Figure 6: Bar chart depicts the distribution of oral lesions based on site. X-axis shows the site of oral mucosal lesions and Y-axis shows the distribution of the study population presented with oral lesions in percentage. Buccal mucosa(63.7%-brown) was the most common site of oral lesions among tobacco users.