

## Effects of Ascorbic Acid Supplement and Furdivita extraction on Hematological Parameters, liver function tests and histopathological changes in Rabbits liver

Qader Abdullah Shannak<sup>1</sup>, Dhamer Ismael Madab<sup>2</sup>, ShifaaAyoob Mohammed<sup>3</sup>,  
Mustafa A. J. Alhiti<sup>4</sup>

<sup>1</sup>MSc. Inorganic Chemistry, In Disease and Histopathological Laboratory, In Biotechnology & Environmental Center, In Fallujah University/Bagdad / Iraq

<sup>2</sup>PhD. Organic Chemistry, Ministry of Education, Salah Al-Din Education Directorate/ Iraq

<sup>3</sup>MSc. Organic Chemistry, Ministry of Education, Salah Al-Din Education Directorate/ Iraq

<sup>4</sup>BSc. Agri. in Pathological Analysis Laboratory, In Biotechnology & Environmental Center, In Fallujah University/Bagdad / Iraq.

\*Corresponding author: [qaderabdullah@uofallujah.edu.iq](mailto:qaderabdullah@uofallujah.edu.iq)

### Abstract

The water-soluble micronutrient Ascorbic (vitamin C) is an essential for several biological activities. Ascorbic acid has a long history of usage in conventional processes that regulate the production of free radicals with antioxidant effects. Free radical levels in cancer cells have been documented to be high. The research aimed at studying hematological of the adult male rabbits. Ascorbic acid (40mg/kg/BW/day for 12 weeks).

The influence of ascorbic acid at the termination of a 12 week clinical cycle on complete counts of erythrocytes (CBR), hemoglobin (HB), cell-packed volume (PCV), total WBC, MCV, MCHC, mean plasma glutathione-s-transferase (GST), acetyl cholinesterase (AChE), and thiobarbituric acid-reactive substances (TBRS). Ascorbic acid therapy resulted in a substantial improvement ( $P < 0.05$ ) in RBC and WBC, while Hb, PCV, MCV, MCH, MCHC, GSH, AChE behavior were not changed in contrast with regulation. Compared to control community, the TBARS are considerably decreased ( $P < 0.05$ ). This search concludes the addition of a vitamin C supply to food will boost anemia and suppress free radicals in adult male rabbits for a long period.

Fordvita is a furan obtained from plant extracts of plant residues *Saccharum officinalis* Effect. This extract was found in the tissues of some organs in mice by watering animals at two concentrations (95.22 and 5.5) mg / ml at LD50 and 50 LD at a rate of one, two, and three weeks. The results showed that both concentrations (95.22 and 5.5) mg / ml have the same thing. The effect on infection compared to the control group. After three weeks, results were better with preserving the organs from histological changes in the intestine, liver and spleen. And that the histological effect was less in the spleen than in the liver and intestine.

The histopathological lesions found by feeding this plant mixture were strengthened in the livers of rabbits with hypercholesterolemia. In this review, diet consumption of the celery plant is recommended; it can be helpful for hypercholesterol and hepatic disease patients.

**Keywords:** Hematological parameters, free radical, acetylcholinesterase and ascorbic acid.

### Introduction

A vitamin C is also available as a dietary substitute and a vitamin present in a range of foods. Vitamin C is also known as ascorbic acid and used for scurvy prevention and recovery. It is often seen as one of the essential vitamins for human and animal health as it tends to repair tissues and create neurotransmitters. Vitamin C is also required for the performance of many

enzymes required for the immune system function. It also works as an antioxidant. [1, 2]. as the recognized name of the vitamin C. It is present in different foods and acts as a nutritional supplement[1]. It is a nutrient. The first vitamin to be chemically developed was vitamin C found in 1912, was extracted in 1928, and in 1933[3]. It is the maximum powerful then safest medicine wanted in a condition system [4].

Vitamin C foods contain citrus, kiwifruit, broccoli, sprouts from Brussels, raw bell peppers, and strawberries[5], respectively. Longer storage or cooking in foods may reduce the amount of vitamin C[5,6]. For normal physiological processes, the body needs vitamin C. It promotes tyrosine, follic acid then tryptophan metabolism[7].

It help to reduce blood saturated fatty acid and to synthesize amino acids that control nervous system carnitine and catecholamine [8]. It is essential to develop tissues and treat wounds. It promotes neurotransmitter formation and improves iron absorption Darkness [9]. It prevents the group from the detrimental impact of the free radical and toxin [6, 11], becoming an antioxidant [10].

Biological pathways, exogenous stimuli and regulation of different enzymes and antioxidant compounds are used to generate free radicals in the body. Oxidative stress arises as the potential of free radical development to defend against it exceeds the potential to avoid it, contributing to trauma, inflammatory actions and chronic situations like atherosclerosis, degenerative diseases and cancer[12].

It has been proposed to mitigate oxidative harm to humans and thus to minimize the likelihood of specific chronic diseases through, vitamin C, vitamin E and  $\beta$ -carotene, also labeled "antioxidant vitamins." In cardiovascular trials, low L-ascorbic acid, tocopherol, and  $\beta$ -carotene plasma are correlated with cardiovascular disease [12, 13]. It can function as an antioxidant in vivo [14, 15] owing to the chemical and biological characteristics of ascorbic acid.

The lack in Vitamin D also has to do with numerous medical conditions such as bone softening, anemia, infections and bleeding gums, scurvy, poor wound healing, haemorrhage, degeneration of organs, atherosclerotic plaques and neurotic disorders. To fix deficiency, vitamin C is also made use of and may not contribute to problems. The research was intended to establish whether vitamin C supplementation has some impact on hematological parameters and amount of serum free radicals in adult male rabbits over a prolonged period of time.

Fordvita extract of amine compounds and nitrofurans which contains an amine group or nitrate at the site [16] of the atom, and it was found that this substance has a positive effect carbon in inhibiting a number of microorganisms. It was also found that it could be furan compounds A therapeutic substance for the treatment of skin infections, the treatment of urinary tract infections and the treatment of infections. In a previous study it was found that Fordvita had no effect on while it had a mixed immunomodulatory effect. [17,18]

Positively, it increased the delay of anaphylaxis, and the effect of the substance on cellular immunity was studied by influencing the phagocytosis process. There was an increase in macrophage efficiency in animals irrigated with Fordvitha extract by measuring the phagocytosis factor. The research targeted the effect of this compound on the tissues of Swiss albino mice after being infected by studying the tissue changes in the organs of these animals[19,20]

### **Material and Method Animals**

Ten adult male (White rabbits) were randomly split into a group of couples and each group had 5 representatives. Community II: Rabbits were provided with vitamin C.

Ascorbic acid continued injected in doses of 40 mg/kg BW once each day for 12 weeks at the end of the time, all rabbits in the community were slaughtered by ether anesthesia. The blood

samples were obtained in different centrifuge tubes. Plasma was isolated by centrifugation at 3000 revolutions a minute for 10 minutes and then rapidly frozen at -20 degrees centigrade for antioxidant enzymes and free radical study.

### Blood Enzyme Activities

The blood sample was collected in two tubes: EDTA (anti-coagulant) and Heparin (anti-coagulant) (anti-coagulant). Non-coagulated blood was tested for total leucocyte numbers (RBC), total erythrocyte counts (WBC), hemoglobin(HB), packed volume of cells (PCV), mean cell volume (MCV), medium hemoglobin (MCH) and mean cell haemoglobin concentration (MCHC) shortly after sampling by the particles counters (from ERMA INC.-Tokio. Model PCE-210) (MCHC). Plasma development activities are computed for Glutathione S-transferase (GST; EC 2.5.1.18) by [21].

The operation of acetylcholinesterase was estimated based on the method was assessed for the plasma-thiobarbituric acid-reactive substances [22].

### Histopathology

Small pieces of liver tissue samples obtained surgically and added to 10% formalin in plastic containers for histopathological examination immediately at room temperature, so sample were fixed by 10% formalin solution for two days then the tissue samples washed up by tap water and processed as the following routine according to [23] for paraffin section and stained with the following haematoxylin and eosin stains.

### Statistical Analysis

Where applicable, statistical analysis was carried out in Minitab software; statistical significance was assessed using two samples T- test analysis after detection normal distribution to the data and appropriate  $P < 0.05$  consider significant [24].

### Results Discussion

As shown in Table1 the data were recorded upon the findings of Ascorbic acid on total erythrocyte counts (RBC), hemoglobin (Hb), packed cell volume (PCV), total leukocyte counts (WBC), mean cell volume (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), plasma glutathione-S-transferease (GST), acetyl cholinesterase (AChE) and thiobarbituric acid-reactive substances (TBARS) at end of the 12-week of treatment.

Treatment with Ascorbic acid produced significant ( $P < 0.05$ ) increase in RBC and WBC, while did not impact any changes in Hb, PCV, MCV, MCH, MCHC, the action of GST and AChE compared to control. TBARS are marginally ( $P < 0.05$ ) decreased compared with control community.

**Table-1: The total means of various parameters at end of 12 week of therapy male rabbits with ascorbic acid (40mg/kg BW)**

Parameters	Animal Groups	
	Control	Ascorbic acid
Total erythrocyte count (RBC; $\times 10^6/\text{mm}^3$ )	5.4± 0.15 <sup>b</sup>	5.8±0.23 <sup>a</sup>
Hemoglobin (Hb; g/dl)	13.9± 0.19 <sup>a</sup>	13.9±0.21 <sup>a</sup>
Packed cell volume (PCV; %)	44.7±0.47 <sup>a</sup>	44.6±0.32 <sup>a</sup>
Total leukocyte count (WBC; $\times 10^3/\text{mm}^3$ )	7.1± 0.51 <sup>b</sup>	7.6±0.48 <sup>a</sup>
Mean cell volume (MCV; fl)	83.2 ± 2.05 <sup>a</sup>	83.4± 2.44 <sup>a</sup>

<b>Mean cell hemoglobin (MCH; pg )</b>	26.5± 0.59 <sup>a</sup>	27.0± 0.59 <sup>a</sup>
<b>Mean cell hemoglobin concentration (MCHC; dl)</b>	31.1± 0.58 <sup>a</sup>	31.3± 0.48 <sup>a</sup>
<b>Glutathione S-transferase (GST; □ mol/hr)</b>	0.88 ± 0.00 <sup>a</sup>	0.90± 0.011 <sup>a</sup>
<b>Acetylcholinesterase (AChE; mol substrate hydrolyzed/ min)</b>	3.3 ± 0.09 <sup>a</sup>	3.3± 0.18 <sup>a</sup>
<b>Thiobarbituric acid-reactive substances (TBARS)</b>	0.30 ± 0.016 <sup>a</sup>	0.23 ± 0.011 <sup>b</sup>

Note: Values are means ± SEM of 5 rabbits in each group. Mean with separate letters (a- b) was slightly difference ( $p \leq 0.05$ ). Mean with the same letters was non-significantly variation ( $p \geq 0.05$ ).

In the present research, a relative increase in BW of 40 mg / kg of ascorbic acid was observed in 12 weeks following previous trials, but the lower dose of ascorbic acid (10 mg / kg / kg BW) was less potent [25]. Hematological parameters such as RBCs, WBCs, MCV, MCH and MCHC are helpful in monitoring the health condition [26].

Present research has demonstrated substantial ( $P < 0.05$ ) rises in RBC and WBC in ascorbic acid therapy, which have not contributed to any Hb, PCV, MCV, MCH or MCHC improvement compared with placebo. The results indicated that treatment with ascorbic acid ameliorated its detrimental effect on hematological parameters.


Similar findings have been reported by another research which study the Impact of ascorbic acid supplementation on the hematological parameters in rabbits [24], in common carp (*Cyprinus carpio* L.) [27] and in Broiler chicks [26]. And this result refers to the absorption of about 80-90 percent ascorbic acid in the gastrointestinal tract.

The ingested acid circulates freely in the sperm, leukocytes and red blood cells, achieving a median plasma concentration of 68-86  $\mu\text{mol/l}$ , with oral intakes of 90-150 mg per day. The body uses it within 2 hours and typically is bloodless in 3 to 4 hours [28]. In detoxifying many chemical substances, GST has been noted as having a significant role [28]. In innervated tissues, AchE is present where its role is to stop the propagation of nerve impulses. The red blood cell membrane also includes it [29].

Compared to control, couple enzymes are not handled ascorbically. These findings have been revised [29]. Free radicals and oxidants are both dangerous and beneficial since they may be hazardous or useful for the body. They are produced either by usual in situ metabolites (pollution, smoke of tobacco, radiation, medication) or by external factors (pollution, smoke of cigarette, radiation, medication) (pollution, cigarette smoke, radiation, medication). When free radicals cannot consistently multiply, they can induce a condition called oxidative stress to concentrate throughout the body.

The human body continuously develops free radical and reactive oxygen species (ROS) [30]. The reactive lipid (LPO) thiobarbituric acid (TBAR) substances are considered to be oxidative stress markers [30]. LPO was assessed by measuring TBARS concentrations in plasma of ascorbic acid processed by male rabbits. The substantial decrease in TBARS in rabbits the treatment with ascorbic acid was in accordance with the previous studies [31, 32]. These experiments were found the rabbits treated with ascorbic acid (20 otherwise 40 mg/kg B.Wt) shows rise in hematological parameters and decrease in TBARS in plasma.


By studying the histological effects. It is observed during the dissection process of mice. For the purpose of taking the organs, apparent changes occurred in the organs of control animals, which were injected at a concentration of 8 CFU / ml.



Picture 1: A cross section of the intestine of a mouse (control) irrigated with physiological solution (40X).


A large spray of inflammatory cells (lymphocytes), plasma cells and neutral cells (degeneration) is observed in the intestine during microscopy. Goblet cells (with expression in Picture 1) were found in comparison with tissues of irrigated animal organs at Fordvita concentrations (95, 22 and 5.5 mg / mL) which showed that Fordvita decreased the rate of infection of animals. The results showed no statistically significant differences (Chi square) in both Fordvita concentrations (5.5 and 95.22) mg / ml.

The tissue sections taken from the liver of mice (control) that were irrigated with physiological solution only and were infected with, changes were observed that included expansion of sinusoids and the occurrence of cavities in different cytoplasm sizes in the cellular plate Increased the delay of anaphylaxis, and the effect of the substance on cellular immunity was studied by influencing the phagocytosis process. There was an increase in macrophage efficiency in animals irrigated with Fordvitha extract



Picture 2: Cross section of the intestine of a mouse irrigated with Fordvita extract (40X).

In the nucleus of hepatic cells, some of them were characterized by high electron density (Pyknotic), while another section was characterized by an increase in the size of the nucleus, and the presence of Histiocytes in the hepatic sinuses and infiltration in the inflammatory cells, especially multicellular PM 3 These changes were not observed when compared with the organ tissues of animals irrigated with Fordvita and both concentrations 5.5 and 95.22 mg / ml and was infected that show in Picture 2.



Picture 3: A cross section of a mouse liver (control) irrigated with physiological solution only and infected with bacteria (40X).




Figure 4: Cross section in the liver of a mouse irrigated with Fordvita extract (40X).

As for the effects caused by Fordvita on the spleen (Spleen), they were less than the effects in other organs. Control, as well as an increase in its inflammatory activity, was observed in the presence of large numbers of cell debris in the cytoplasm (Figure 3,4). Also, no significant differences were observed (square chi) in both concentrations of Fordvita 5.5 and 95.22 mg / mL.



Picture 5: A section of the spleen of a mouse (control) irrigated with physiological solution only (40X).

The effect in all the members after three weeks was better than the first and second weeks, due to the absence of any tissue changes. Infection. Has become widespread all over the world, and its epidemic is high in hospitals with increasing antibiotic resistance and the

importance of using nitrofuran compounds as a drug against infection. Is one of the most important methods of controlling the disease (33-39).




Figure 6: A section in the spleen of a mouse irrigated with Fordvita extract (40X).

In other studies, the number of white blood cells (WBC) was calculated in cases of Gram-negative bacteria. In rabbits, to find out the state of Neutropenia and their results showed the benefit of the effect of treatment with furan derivatives, but only at the onset of infection with Gram-negative bacteria. In the tissues after intravenous injection of nitrofuran compounds (39-44).

The results that appeared in this study are similar to those reached by previous researchers in extending the efficacy of furans and nitrofuran derivatives in treating infection, especially the compound vancomycine, as well as the treatment effect on the liver and intestine when compared with the spleen in Figure 5,6 . While it differed from another study, it mentioned the inefficiency of treatment with nitrofuran compounds in the treatment of germ-positive bacteria, such as *Staphylococcus aureus*. (45,46).

### Conclusions

Ascorbic acid is an antioxidant that preserves the body from the harm incurred by free radicals. It is seen in a number of health ailments. The ascorbic acid concentration of 40 mg kg was found to be optimal to boost the hematological parameters and decrease the free radical level in male rabbits in an experiment.

### REFERENCES

1. **Angulo, C., Maldonado, R., Pulgar, E., Mancilla, H., Córdova, A., Villarroel, F., & Concha, I. I. (2011).** Vitamin C and oxidative stress in the seminiferous epithelium. *Biological research*, 44(2), 169-180.
2. **Chambial, S., Dwivedi, S., Shukla, K. K., John, P. J., & Sharma, P. (2013).** Vitamin C in disease prevention and cure: an overview. *Indian Journal of Clinical Biochemistry*, 28(4), 314-328.
3. **Daniel, E. P., & Munsell, H. E. (1937).** Vitamin content of foods: a summary of the chemistry of vitamins, units of measurement, quantitative aspects in human nutrition and occurrence in foods (No. 275). US Department of Agriculture.
4. **Dave, K. N., & Patil, R. S. (2017).** Biological Importance of Ascorbic Acid (Vitamin C) in Human Health-A Classic Review. *International Journal For Research In Biology & Pharmacy* (ISSN: 2208-2093), 3(7), 1-8.
5. **El-Demerdash, F. M., Yousef, M. I., Kedwany, F. S., & Baghdadi, H. H.**

- (2004). Cadmium-induced changes in lipid peroxidation, blood hematology, biochemical parameters and semen quality of male rats: protective role of vitamin E and  $\beta$ -carotene. *Food and chemical toxicology*, 42(10), 1563-1571.
6. **Ellman, G. L., Courtney, K. D., Andres Jr, V., & Featherstone, R. M. (1961).** A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical pharmacology*, 7(2), 88-95.
  7. **Habig, W. H., Pabst, M. J., & Jakoby, W. B. (1974).** Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *Journal of biological Chemistry*, 249(22), 7130-7139.
  8. **Iqbal, K., Khan, A., & Khattak, M. M. A. K. (2004).** Biological significance of ascorbic acid (vitamin C) in human health-a review. *Pakistan Journal of Nutrition*, 3(1), 5-13.
  9. **Johnson, L. J., Meacham, S. L., & Kruskall, L. J. (2003).** The antioxidants-vitamin C, vitamin E, selenium, and carotenoids. *Journal of agromedicine*, 9(1), 65-82.
  10. **Khan, P. K., & Sinha, S. P. (1996).** Ameliorating effect of vitamin C on murine sperm toxicity induced by three pesticides (endosulfan, phosphamidon and mancozeb). *Mutagenesis*, 11(1), 33-36.
  11. **Monsen, E. R. (2000).** Dietary reference intakes for the antioxidant nutrients: vitamin C, vitamin E, selenium, and carotenoids. *Journal of the Academy of Nutrition and Dietetics*, 100(6), 637.
  12. **Ogunlesi, M., Okiei, W., Azeez, L., Obakachi, V., Osunsanmi, M., & Nkenchor, G. (2010).** Vitamin C contents of tropical vegetables and foods determined by voltammetric and titrimetric methods and their relevance to the medicinal uses of the plants. *Int J Electrochem Sci*, 5, 105-115.
  13. **Olson, J. A., (1987).** Recommended dietary intakes (RDI) of vitamin A in humans. *The American journal of clinical nutrition*, 45(4), 704-716.
  14. **Oyawoye, B. M., & Ogunkunle, H. N. (2004).** Biochemical and haematological reference values in normal experimental animals. *New York: Mason*, 212-216.
  15. **Paulson, D. S. (2008).** Biostatistics and microbiology: a survival manual. *Springer Science & Business Media*.
  16. **Pham-Huy, L. A., He, H., & Pham-Huy, C. (2008).** Free radicals, antioxidants in disease and health. *International journal of biomedical science: IJBS*, 4(2), 89.
  17. **Rock, C. L., Jacob, R. A., & Bowen, P. E. (1996).** Update on the biological characteristics of the antioxidant micronutrients: vitamin C, vitamin E, and the carotenoids. *Journal of the American Dietetic Association*, 96(7), 693-702.
  18. **Rose, R. C., & Bode, A. M. (1993).** Biology of free radical scavengers: an evaluation of ascorbate. *The FASEB journal*, 7(12), 1135-1142.
  19. **Sato, K., Niki, E., & Shimasaki, H. (1990).** Free radical-mediated chain oxidation of low density lipoprotein and its synergistic inhibition by vitamin E and vitamin C. *Archives of Biochemistry and Biophysics*, 279(2), 402-405.
  20. **Semba, R. D. (2012).** The discovery of the vitamins. *International Journal of Vitamin and Nutrition Research*, 82(5), 310-315.
  21. **Sharaf, R., Khan, A., Khan, M. Z., Hussain, I., Abbas, R. Z., Gul, S. T., ...& Saleemi, M. K. (2013).** Arsenic Induced Toxicity in Broiler Chicks and Its Amelioration with Ascorbic Acid: Clinical, Hematological and Pathological Study. *Pakistan Veterinary Journal*, 33(3).
  22. **Souza, M. S., Magnarelli, G. G., Rovedatti, M. G., Cruz, S. S., & D'Angelo, A. P. D. (2005).** Prenatal exposure to pesticides: analysis of human placental acetylcholinesterase, glutathione S-transferase and catalase as biomarkers of effect.



- Biomarkers, 10(5), 376-389.
23. **Subasree, S. (2014).** Role of vitamin C and vitamin E in health and disease. *Journal of Pharmaceutical Sciences and Research*, 6(1), 52.
  24. **Tappel, A. L., & Zalkin, H. (1959).** Inhibition of lipide peroxidation in mitochondria by vitamin E. *Archives of Biochemistry and Biophysics*, 80(2), 333-336.
  25. **Villacorta, L., Azzi, A., & Zingg, J. M. (2007).** Regulatory role of vitamins E and C on extracellular matrix components of the vascular system. *Molecular aspects of medicine*, 28(5-6), 507-537.
  26. **Vinodhini, R., & Narayanan, M. (2009).** The impact of toxic heavy metals on the hematological parameters in common carp (*Cyprinus carpio* L.). *Journal of Environmental Health Science & Engineering*, 6(1), 23-28.
  27. **World Health Organization (WHO). (2016).** WHO Model List of Essential Medicines-19th List. April 2015-Amended November 2015. Available at: [who.int/medicines/publications/essentialmedicines/en/](http://who.int/medicines/publications/essentialmedicines/en/). Accessed November, 2.
  28. **Yousef, M. I., Abdallah, G. A., & Kamel, K. I. (2003).** Effect of ascorbic acid and vitamin E supplementation on semen quality and biochemical parameters of male rabbits. *Animal Reproduction Science*, 76(1-2), 99-111.
  29. **Yousef, M. I., El-Morsy, A. M., & Hassan, M. S. (2005).** Aluminium-induced deterioration in reproductive performance and seminal plasma biochemistry of male rabbits: protective role of ascorbic acid. *Toxicology*, 215(1-2), 97-107.
  30. **Yousef, M. I., Salem, M. H., Kamel, K. I., Hassan, G. A., & El-Nouty, F. D. (2003).** Influence of ascorbic acid supplementation on the haematological and clinical biochemistry parameters of male rabbits exposed to aflatoxin B1. *Journal of Environmental Science and Health, Part B*, 38(2), 193-209.
  31. **Yousef, M. I., Salem, M. H., Kamel, K. I., Hassan, G. A., & El-Nouty, F. D. (2003).** Influence of ascorbic acid supplementation on the haematological and clinical biochemistry parameters of male rabbits exposed to aflatoxin B1. *Journal of Environmental Science and Health, Part B*, 38(2), 193-209.
  32. **ALP, E; Gozukucuk, S; Canoz, O; Kirmaci, B; And Doganay, M 2004,** Effect of granulocyte colony – stimulating factor in experimental methicillin resistant, *BMC infect.Dis.* 4:43.
  33. **Al-Saloom, R.H.; Al-Hamendo, M.M.; and Abu-Risha, R.A. 2005.** Study the effect of Furdivita extract on cellular and humoral immunity. *Iraqi J. Biotech. res.* 7(1): 18-25
  34. **Craignill, L. 1992.** Update to Nitrofurans. *Environ. Toxicol.* 12 (2).
  35. **Culling, C.F.A 1981.** Handbook of histopathological and histochemical techniques, 3 rd. Butterwoth. P: 712.
  36. **Deshazo, R.D. and Kemp; S. F. 1997** Allergic reaction to drug and biological agents. *JAMA.* 22, 1895-1906.
  37. **Humson, C.L 1972.** In: animal tissue techniques (3rd Ed.). P: 41, W.H. Freeman and company.
  38. **Lautenschlager, S; Herzog, C; and Zimmerli, W. 1993.** Course and outcome of bacteremia due to *S.aureus*: evaluation of different clinical case definitions. *Clin.infect.Dis.* 16:567-573.
  39. **Marion, I. and Ias. 1976.** Absorption of nitrofurazon *Curr. The Res.* 19: 550.
  40. **Panlilio, A.L; culver, D.H; Gaynes, R.P; Banerjee, S; Henderson, T.S; Tolson, J.S; and Marton, W.J. 1992.** Methicillin resistant *S. aureus* in US hospital, 1975-1991. *Infect. Control hosp. Epidmiol.* 13: 582-586.

41. **Sieradzki, K.; Roberts, R.B., Haber, S.W; and Tomasz, A. 1999.** The development of vancomycin Resistance in a patient with methicillin. Resistant Staphylococcus aureus infection. The new English J.Med.340 (7):517 – 523.
42. **Smith, W.S; Sumnicht, G.E ; Sharpe, R.W; Samuelson, S.D; and Nillard, F.E.1995.** GCSF versus placebo in addition to penicillin – G in a randomized blinded study of gram – negative pneumonia sepsis : analysis of survival and multisystem organ failure . Blood. 86: 1301 – 1309.
43. **Tilles, S.A. 2001.** Patricate Issues in the Management of types reactions. Sulfonamides. South. Med. 94 (8).
44. **Wolf, M.S. 1975** furazolidone antibiotic used against Giardiasis. J. Am. Med. Ass. 233: 1362.
45. **Yamauchi, T; Ishibashi, A.; Shikata, K.; Tokuhara, N.; Seino, K. Kobayashi; S; and Nagai, M. 2004.** Effect of E6060 ( 4 – 5 – 7 fluoro – 4 (trifluoromethyl ) benzo ( b ) Furan – 2 – YL )-1H-2- pyrrolyl)benzoic acid) a novel subtype-selective retinoid on lupus – like nephritis in female J.pharm and Exerim .Therap , Fast Forward , 10: 1124 .
46. **Yao, L; Berman, J.W; Factor, S.M; and Lowy, F.D. 1997.** Correlation of histopathologic and bacteriologic changes with cytokine expression in an experimental murine model of bacteremic S.aureus infection. Infect.Immun.65:3889- 3895.