

Study on Fuzzy Game Problem in Icosikaitetragonal Fuzzy Number

S. Sasikumar¹ and V. Raju²

¹Research Scholar, Department of Mathematics,
Vels Institute of Science, Technology and Advanced Studies (VISTAS),
Chennai-600 117, Tamil Nadu, India

² Assistant Professor, Department of Mathematics
School of Basic Sciences
Vels Institute of Science, Technology and Advanced Studies (VISTAS),
Chennai-600 117, India

¹ sasikumar_math@yahoo.co.in and ² matraju@gmail.com

ABSTRACT

In this article, we bring in the fuzzy game problem by Icosikaitetragonal fuzzy number. We generate the value of Payoff matrix by Icosikaitetragonal fuzzy number. We modify the fuzzy game problem into crisp valued game problem by using ranking to pay off. The crisp valued game problem can be solved by Oddment method and illustrations are exemplified

Keywords

Icosikaitetragonal fuzzy number, fuzzy game problem, fuzzy ranking, pay off matrix.

1.Introduction

Fuzzy set theory was introduced by Zadeh [1].The concept of fuzzy set theory reveals imprecision and vagueness. We make use of the fuzzy set almost in all business and also in our many day to day life activities. We bring together more information from the environment in each day is fuzzy. We are not able to move our daily life without crossing fuzzy connected circumstance. Fuzzy set is efficient in many ongoing world situations. Jain [2] was the first to propose method of ranking fuzzy numbers for decision making in fuzzy related situation. Raju and Jayagopal [3] was the first to introduce the Icosikaitetragonal fuzzy number. Game theory makes it easier to have interaction of decision makers between the scenario of co operation and spirited approach.

Game theory has extensive range of applications in various fields such as business model development, Economics, diplomacy and military strategy. In a game theory each player is to take good decision by selecting various strategies from the all available strategies. When uncertainties occur in game theory, fuzzy set is the best tool to study this kind of game in which pay off matrices is denoted by fuzzy numbers. In this paper, we have taken two person zero sum game, in which imprecise values are Icosikaitetragonal fuzzy numbers. We have made clear it with converting to crisp valued game problem using ranking technique. We have analyzed fuzzy game problem using Icosikaitetragonal fuzzy number with illustrations and solved the game problem with oddment method

2. PRELIMINARIES

In this section, we give the preliminaries that are required for this study.

Definition 2.1. A fuzzy set A is defined by $A = \{(x, \mu_A(x)) : x \in A, \mu_A(x) \in [0, 1]\}$. Here x is crisp set A and $\mu_A(x)$ is membership function in the interval $[0, 1]$.

Definition 2.2.

The fuzzy number A is a fuzzy set whose membership function must satisfy the following conditions.

- (i) A fuzzy set A of the universe of discourse X is convex
- (ii) A fuzzy set A of the universe of discourse X is a normal fuzzy set if $x_i \in X$ exists
- (iii) $\mu_A(x)$ is piecewise continuous

Definition 2.3.

A fuzzy number $A = (a, b, c)$, where $a \leq b \leq c$, is triangular fuzzy number and its membership function is given by

$$\mu_A(x) = \begin{cases} \frac{x-a}{b-a}, & \text{for } a \leq x \leq b \\ \frac{c-x}{c-b}, & \text{for } b \leq x \leq c \\ 0, & x > c \end{cases}$$

Definition 2.4

A fuzzy number $A = (a, b, c, d)$, where $a \leq b \leq c \leq d$, is trapezoidal fuzzy number and its membership function is given by

$$\mu_A(x) = \begin{cases} 0, & \text{for } x < a \\ \frac{x-a}{b-a}, & \text{for } a \leq x \leq b \\ 1, & \text{for } b \leq x \leq c \\ \frac{d-x}{d-c}, & \text{for } c \leq x \leq d \\ 0, & x > d \end{cases}$$

Definition 2.5

An α -cut of fuzzy set A is classical set defined as ${}^\alpha A = \{x \in X | \mu_A(x) \geq \alpha\}$

Definition 2.6

A fuzzy set A is a convex fuzzy set iff each of its α -cut ${}^\alpha A$ is a convex set.

Definition 2.7

Game theory provides a mathematical framework for analyzing the decision-making processes and strategies of adversaries (or *players*) in different types of competitive situations. The simplest type of competitive situations is two-person, zero-sum games. These games involve only two players; they are called *zero-sum* games because one player wins whatever the other player loses.

Definition 2.8 [3] A fuzzy number $A = (a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, \dots, a_{24})$ is Icosikaitetragonal fuzzy number and its membership function is given by

$$\mu_A(x) = \begin{cases} 0, & \text{for } x < a_1 \\ k_1 \left(\frac{x - a_1}{a_2 - a_1} \right), & \text{for } a_1 \leq x \leq a_2 \\ k_1, & \text{for } a_2 \leq x \leq a_3 \\ k_1 + (k_2 - k_1) \left(\frac{x - a_3}{a_4 - a_3} \right), & \text{for } a_3 \leq x \leq a_4 \\ k_2, & \text{for } a_4 \leq x \leq a_5 \\ k_2 + (k_3 - k_2) \left(\frac{x - a_5}{a_6 - a_5} \right), & \text{for } a_5 \leq x \leq a_6 \\ k_3, & \text{for } a_6 \leq x \leq a_7 \\ k_3 + (k_4 - k_3) \left(\frac{x - a_7}{a_8 - a_7} \right), & \text{for } a_7 \leq x \leq a_8 \\ k_4, & \text{for } a_8 \leq x \leq a_9 \\ k_4 + (k_5 - k_4) \left(\frac{x - a_9}{a_{10} - a_9} \right), & \text{for } a_9 \leq x \leq a_{10} \\ k_5, & \text{for } a_{10} \leq x \leq a_{11} \\ k_5 + (1 - k_5) \left(\frac{x - a_{11}}{a_{12} - a_{11}} \right), & \text{for } a_{11} \leq x \leq a_{12} \\ 1, & \text{for } a_{12} \leq x \leq a_{13} \\ k_5 + (1 - k_5) \left(\frac{a_{14} - x}{a_{14} - a_{13}} \right), & \text{for } a_{13} \leq x \leq a_{14} \\ k_5, & \text{for } a_{14} \leq x \leq a_{15} \\ k_4 + (k_5 - k_4) \left(\frac{a_{16} - x}{a_{16} - a_{15}} \right), & \text{for } a_{15} \leq x \leq a_{16} \\ k_4, & \text{for } a_{16} \leq x \leq a_{17} \\ k_3 + (k_4 - k_3) \left(\frac{a_{18} - x}{a_{18} - a_{17}} \right), & \text{for } a_{17} \leq x \leq a_{18} \\ k_3, & \text{for } a_{18} \leq x \leq a_{19} \\ k_2 + (k_3 - k_2) \left(\frac{a_{20} - x}{a_{20} - a_{19}} \right), & \text{for } a_{19} \leq x \leq a_{20} \\ k_2, & \text{for } a_{20} \leq x \leq a_{21} \\ k_1 + (k_2 - k_1) \left(\frac{a_{22} - x}{a_{22} - a_{21}} \right), & \text{for } a_{21} \leq x \leq a_{22} \\ k_1, & \text{for } a_{22} \leq x \leq a_{23} \\ k_1 \left(\frac{a_{24} - x}{a_{24} - a_{23}} \right), & \text{for } a_{23} \leq x \leq a_{24} \\ 0, & \text{for } x > a_{24} \end{cases}$$

3. Mathematical formulation of Fuzzy Game problem:

Consider a two person zero sum game in which all the entries of the payoff matrix are Icosikaitetragonal fuzzy numbers. Let us take the player P has m strategies and player Q has n strategies. We will assume that each player has to choose from the pure strategies. Player P is always considered to be gainer and the player Q is always loser. Then the payoff matrix m x n is

$$G = \begin{pmatrix} p_{11} & p_{12} & \cdot & p_{1n} \\ p_{21} & p_{22} & \cdot & p_{2n} \\ \cdot & \cdot & \cdot & \cdot \\ p_{m1} & p_{m2} & \cdot & p_{mn} \end{pmatrix}$$

3.1 Numerical Examples:

Consider the fuzzy game problem with payoff matrix as Icosikaitetragonal fuzzy numbers. This problem is worked out by taking the values

$$k_1 = \frac{1}{6}, k_2 = \frac{2}{6}, k_3 = \frac{3}{6}, k_4 = \frac{4}{6}, k_5 = \frac{5}{6}$$

We get the values of $\mu_{Icsktetra}(a_{ij})$

a_{11}	-13,-12,-11,-10,-9,-8,-7,-6,-4,-3,-2,0,1,2,4,6,8,10,12,14,16,18,20,22	$\mu_{Icsktetra}(a_{11}) = 2$
a_{12}	1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24	$\mu_{Icsktetra}(a_{12}) = 12.5$
a_{21}	-12,-11,-10,-9,-8,-7,-6,-5,-4,-3,-2,-1,0,3,5,6,7,8,10,12,14,15,16,18	$\mu_{Icsktetra}(a_{21}) = 1.67$
a_{22}	2,4,6,8,10,12,14,16,18,20,22,24,26,28,30,32,34,36,38,40,42,44,46,48	$\mu_{Icsktetra}(a_{22}) = 25$
a_{31}	0,1,2,3,4,5,6,7,8,9,10,11,13,15,17,19,21,23,25,27,29,30,31,33	$\mu_{Icsktetra}(a_{31}) = 14.54$
a_{32}	1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33,35,37,39,41,43,45,47	$\mu_{Icsktetra}(a_{32}) = 24$

Fuzzy game problem is reduced to the following payoff matrix $A = \begin{pmatrix} 2 & 12.5 \\ 1.67 & 25 \\ 14.54 & 24 \end{pmatrix}$

Minimum of first row is 2

Minimum of second row is 1.67

Minimum of third row is 14.54

Maximum of first column is 14.54

Maximum of second column is 25


Max(min) = 14.54

Min(max) = 14.54

It has a saddle point

Value of the game is 14.54

3.2 Diagram of Icosikaitetragonal fuzzy number:



3.3 Ranking of Icosikaitetragonal fuzzy number:

Let I be a normal Icosikaitetragonal fuzzy number. The value $M(I)$, called as measure of I is calculated as

$$M(I) = \frac{1}{2} \int_1^{k_1} (\ell_1 + \ell_2) d\ell + \frac{1}{2} \int_{k_1}^{k_2} (m_1 + m_2) dm + \int_{k_2}^{k_3} (n_1 + n_2) dn + \int_{k_3}^{k_4} (o_1 + o_2) do + \int_{k_4}^{k_5} (p_1 + p_2) dp + \int_{k_5}^1 (q_1 + q_2) dq$$

where $0 \leq k_1 \leq k_2 \leq k_3 \leq k_4 \leq k_5 \leq 1$

$$M(L) = \frac{1}{4} \left[(a_1 + a_2 + a_{23} + a_{24})k_1 + (a_3 + a_4 + a_{21} + a_{22})(k_2 - k_1) + (a_5 + a_6 + a_{19} + a_{20})(k_3 - k_2) + (a_7 + a_8 + a_{17} + a_{18})(k_4 - k_3) + (a_9 + a_{10} + a_{15} + a_{16})(k_5 - k_4) + (a_{11} + a_{12} + a_{13} + a_{14})(1 - k_5) \right]$$

where $0 \leq k_1 \leq k_2 \leq k_3 \leq k_4 \leq k_5 \leq 1$

we take the values for $k_1 = \frac{1}{6}, k_2 = \frac{2}{6}, k_3 = \frac{3}{6}, k_4 = \frac{4}{6}, k_5 = \frac{5}{6}$

3.4 Numerical Examples:

Let us consider the matrix

$$\begin{pmatrix} (-6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17) & (-8, -7, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15) & (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24) \\ (-6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18) & (-4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19) & (0, 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 20, 22, 24, 26, 27, 28, 29, 30) \\ (0, 1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 19, 21, 22, 24, 25, 26, 27, 28, 29) & (1, 2, 3, 6, 8, 9, 10, 12, 13, 15, 16, 17, 19, 20, 22, 23, 25, 28, 30, 32, 34, 35, 37, 39) & (-5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18) \end{pmatrix}$$

Step 1:

We obtain the values of $\mu_{Icskoc}(a_{ij})$ of the given fuzzy game problem and convert the fuzzy game problem into crisp valued problem which is shown in the given table.

a_{11}	$-6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17$	$\mu_R(a_{11}) = 5.5$
a_{12}	$-8, -7, -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15$	$\mu_R(a_{12}) = 3.5$
a_{13}	$1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24$	$\mu_R(a_{13}) = 12.5$
a_{21}	$-6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18$	$\mu_R(a_{21}) = 5.54$
a_{22}	$-4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19$	$\mu_R(a_{22}) = 7.5$
a_{23}	$0, 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 18, 20, 22, 24, 26, 27, 28, 29, 30$	$\mu_R(a_{23}) = 14.3$
a_{31}	$0, 1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 19, 21, 22, 24, 25, 26, 27, 28, 29$	$\mu_R(a_{31}) = 14.25$
a_{32}	$1, 2, 3, 6, 8, 9, 10, 12, 13, 15, 16, 17, 19, 20, 22, 23, 25, 28, 30, 32, 34, 35, 37, 39$	$\mu_R(a_{32}) = 19$
a_{33}	$-5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18$	$\mu_R(a_{33}) = 6.5$

Step 2 : The given fuzzy game problem is reduced to the following payoff matrix

$$\begin{matrix} & \mathbf{B} \\ \mathbf{A} & \begin{pmatrix} 5.5 & 3.5 & 12.5 \\ 5.54 & 7.5 & 14.3 \\ 14.25 & 19 & 6.5 \end{pmatrix} \end{matrix}$$

Minimum of first row = 3.5

Minimum of second row = 5.54

Minimum of third row = 6.5

Maximum of first column = 14.25

Maximum of second column = 19

Maximum of third column = 14.3

Max (min) = 6.5

Min (max) = 14.25

Max (min) $\neq$ Min (max)

6.5 $\neq$ 14.5

There is no saddle point

Step 3: To solve the value of the problem, we apply dominance method. Clearly first row is dominated by second row as all the elements of first row are less than second row. Hence eliminate first row, we get

$$\begin{matrix} & \mathbf{B} \\ \mathbf{A} & \begin{pmatrix} 5.54 & 7.5 & 14.3 \\ 14.25 & 19 & 6.5 \end{pmatrix} \end{matrix}$$

Again first column is dominated by second column. So eliminate first column.

$$\begin{matrix} & \mathbf{B} \\ \mathbf{A} & \begin{pmatrix} 7.5 & 14.3 \\ 19 & 6.5 \end{pmatrix} \end{matrix}$$

Now we obtain 2x2 payoff matrix. To solve the reduced matrix, which won't have any saddle point. So we apply oddment method. Hence the augmented payoff matrix is

$$\begin{matrix} & \mathbf{B} \\ \mathbf{A} & \begin{pmatrix} 7.5 & 14.3 \\ 19 & 6.5 \end{pmatrix} \end{matrix}$$

$$(7.5P_1) + 14.3(1 - P_1) = 19P_1 + 6.5(1 - P_1)$$

$$P_1 = 0.4041$$

$$1- P_1 = 0.5959$$

$$7.5 Q_1 + 19 (1- Q_1) = 14.3 Q_1 + 6.5(1- Q_1)$$

$$Q_1 = 0.6477$$

$$1- Q_1 = 0.3523$$

$$\text{Strategy for B} = (0.4041, 0, 0.5959)$$

$$\text{Strategy for A} = (0.6477, 0, 0.3523)$$

$$\text{Value of the game} = (7.5 P_1) + 14.3(1- P_1)$$

$$= (7.5 \times 0.4041) + (14.3 \times 0.5959) = 11.55212$$

Conclusion: In this article, we have examined and solved 3×3 fuzzy pay off matrix whose elements are Icosikaitetragonal fuzzy number. We have illustrated the optimal solution of the fuzzy valued game problem converting to crisp valued game problem using ranking techniques. The Crisp valued game problem is solved by oddment method

References:

- [1] L.A. Zadeh, , Fuzzy sets, Information and Control, 8(3) ,1965, 338-353.
- [2] R.E.Bellman and L.A.Zadeh, Decision making in fuzzy environment, Management Science, 17, 1970, 141- 164.
- [3] V. Raju and R. Jayagopal, “A new operation on Icosikaitetragonal fuzzy number”, Journal of Combinatorial Mathematics and Combinatorial Computing ,Volume 112(2020), Page no : 127-136
- [4] V. Raju and R. Jayagopal “An Arithmetic Operations of Icosagonal fuzzy number using Alpha cut ”International Journal of Pure and Applied Mathematics. Volume 120, No. 8, 2018, 137-145
- [5] V. Raju and R. Jayagopal “An Approach on Icosikaioctagonal Fuzzy number- Traditional Operations on Icosikaioctagonal fuzzy number” A Journal of composition theory, Vol.XII, Issue X, 2019, Page No: 727-734
- [6] V. Raju and R. Jayagopal “ A Rudimentary Operations on Octagonal Fuzzy Numbers ” International Journal of Research in Advent Technology Vol.6, No.6, June 2018, Page No: 1320-1323
- [7] V.Raju and M.ParuvathaVathana “Discourse on Fuzzy Game Problem in Icosagonal Fuzzy Number ” International journal of scientific research and review volume 8, Issue 3, 2019, Page no: 1384-1390
- [8] V.Raju and S.Maria Jesu Raja “An Approach on Fuzzy game problem in Icosikaioctagonal Fuzzy number” Journal of Xidian University, Volume 14, Issue 4, 2020, Page no: 1009-1016
- [9] V.Raju , Ranking Function on Icosagonal Fuzzy Number for Solving Fuzzy Transportation Problem, “Journal of Applied Science and Computations ” Volume VI, Issue IV, 2019, Page No: 3631-3640

- [10] V.Raju and M.ParuvathaVathana “An Icosagonal Fuzzy Number for solving Fuzzy Sequence Problem” International journal of Research in Engineering, IT and Social Sciences” Volume 9, Issue 5, 2019. Page no: 37-40
- [11] V.Raju and M.ParuvathaVathana “ Fuzzy Critical path method with Icosagonal Fuzzy Numbers using Ranking Method” A Journal of Composition Theory, Volume 12, Issue 9, 2019, Page no: 62-69
- [12] V.Raju and S.Arul Amirtha Raja “ Study on fuzzy sequencing problem in Icosikaioctagonal Fuzzy Numbers” Journal of Xidian University, Volume 14, Issue 4, 2020, Page no: 3829-3837
- [13] V.Raju and S.Maria Jesu Raja “ Fuzzy decision making problem in Icosikaioctagonal Fuzzy number” Journal of Xidian University, Volume 14, Issue 5, 2020, Page no: 3240-3248
- [14] K. Arulmozhi , V. Chinnadurai “Bipolar fuzzy soft hyper ideals of ordered –hypersemigroups Γ ” International Journal of Scientific Research and Review Volume 8, Issue 1, 2019, Page No: 1134-1140