

Bisphenol A (BPA) as an Endocrine Disrupting Chemical (EDC) -A Review

Chanemougavally. J¹, Thotakura Balaji², Shruthy.K.M³, Prabhu. K⁴

¹Tutor, Department of Anatomy, A.C.S Medical College and Hospital, Dr. M.G.R Educational and Research Institute (Deemed to be University), Velappanchavadi, Chennai- 600077, Tamilnadu, India.

²Professor, Department of Anatomy, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education (Deemed to be University), Kelambakkam, Chennai- 603103. Tamilnadu, India.

³Associate Professor, Department of Anatomy, A.C.S Medical College and Hospital, Dr. M.G.R Educational and Research Institute (Deemed to be University), Velappanchavadi, Chennai- 600077, Tamilnadu, India.

⁴Associate Professor, Department of Anatomy, Shree Balaji Medical College and Hospitals, Bharath Institute of Higher Education and Research (Deemed to be University) Chromepet, Chennai- 600044, Tamilnadu, India.

Name and Postal Address of the corresponding author:

J. Chanemougavally, M.Sc., (Anatomy)

Department of Anatomy, A.C.S Medical College and Hospitals,

Dr. MGR Educational and Research Institute (Deemed to be University),

Velappanchavadi, Chennai – 600077.

Tamilnadu, India.

e-mail: chanemou@gmail.com

E-mail address for all contributing authors:

1.Chanemougavally. J chanemou@gmail.com

2. Thotakura Balaji thotakurabalaji@gmail.com

3. Shruthy K.M shrumukundan@gmail.com

4. Prabhu. K prbanu75@gmail.com

Abstract

BPA stands for Bisphenol A [22 – Bis (4 - Hydroxyphenol) propane] is an organic synthetic compound, most commonly exposed chemical in everyday life. BPA is used in hard plastic bottles and as a protective lining in canned food and beverage cans. Bisphenol A (BPA) is an old synthetic chemical widely known for endocrine malfunctions and replacement of the estrogenic activity of diethylstilbestrol.¹ Various environmental pollutants are manmade chemicals like polymerised form of biphenyls and bisphenol-A have several developmental effects on endocrine and reproductive functions of living beings. Bisphenol-A

causes wider effects even at low doses in both terrestrial and aquatic animals and effects major disturbances at endocrine, immune and metabolic level.² Since BPA is known to have 'selective estrogen receptor modulator' properties and epigenetic regulations. It affects multiple reproductive organs such as testis, ovaries etc. as it causes reproductive biohazard and endocrine disruption.³ This review summarizes the highlights of BPA as an Endocrine disrupting chemical (EDC) and how it effects multiple organs in the body.

Key Words: Bisphenol A, Endocrine Disruptor, Estrogen

Introduction

Bisphenol A (BPA) is an estrogenic chemical which has been used widely in the production of polycarbonate plastics and epoxy resins.⁴ It was **first synthesized in 1891 by Aleksandr** (Russia) and first mention of BPA in scientific paper was made in 1905 by Thomas Zincke (Germany). Many household electronic appliances, several medical devices, safety equipment used in sports, plumbing equipment are few objects that contain BPA. BPA is also used as a supplement in the production of brominated flame retardants and brake fluid, Bisphenol Aglycidyl methacrylate and bisphenol A-dimethacrylate are BPA derivatives used as a component in the production of dental fillings and sealants in the dental industry.¹ With BPA being used so widely, it has drawn immense attention due to its various effects on human health⁵.

BPA detected in canned and brewed food cause health problems especially for developing foetus, infants and children. The estimated exposure levels of BPA are from **0.01 µg/kg/day in Children** and **4.2 µg/kg/day in adults**. BPA consumption causes male infertility, heart diseases, depression, Type 2 Diabetes, loss of memory, asthma and carcinoma of brain. BPA being a potent reproductive and genotoxic agent has severe effects on the normal physiological functions⁴. BPA has a weak estrogenic activity and can interfere with the endocrine system, and can increase the possibility of testis and breast cancer⁶.

The major adverse effects of Endocrine Disrupting Chemicals (EDC) are observed in hormone system malfunction. It alters the essential physiological functions by distorting endocrine regulation. There are potent EDCs in the environment such as natural and man-made steroid estrogens which includes 17β-estradiol (E2), estrone (E1), and 17 α-ethinylestradiol (EE2). Bisphenol A (BPA) and nonylphenol (NP) are other common industrial chemicals involved in interfering the function of endocrine system in animals⁷.

Endocrine Disrupting Chemical (EDC)

EDCs are exogenous agents that interfere with the function of hormones like secretion, release, transport, binding, action and metabolism that are responsible for the maintenance of self-equilibrium and the regulation of developmental processes⁷. Natural and synthetic chemicals interfere with the body's hormones (endocrine system). EDCs have effects on the normal organ development, reproductive system, brain function, immune system as the Endocrine disruptors are found in several routine products such as food

containers, plastic water bottles, hot and cold metal soft drink cans, detergents, baby toys, wider range of cosmetics and pesticides⁸.

Source of EDC

There are multiple routes of entry for EDCs into the environment from diverse sources. It can be from the air, soil or water. EDC sediments have been traced in both in biota and fish that enter the aquatic ecosystem through wastewater, agricultural run-off and groundwater discharge. Significant contributors of EDCs to surface waters are discharges from municipal wastewater treatment plants (WWTPs)⁷.

Discharges to water treatment facilities include hormones, pharmaceutical estrogens excreted by humans flushed down from home toilets, PPCPs excreted or washed from the body, plant material, waste products from fire retardants, regular household cleaning products, insecticides and pesticides are the main sources. WWTPs might also receive waste products from industrial processes that use plastics containing BPAs and storm water run-off that contain EDCs.⁷

Endocrine-disrupting chemicals usually takes time to break-down in the environment. Thus, makes them potentially hazardous over time. Endocrine disrupting chemicals also causes adverse effects in animals. Human beings are also exposed to EDC, however to assess its effects on health is an enormous task. Moreover, traces of endocrine-disrupting chemicals may be unsafe. Normal endocrine functioning in the body involves very minimal changes in hormone levels. However, these small fluctuations are sufficient to bring out significant developmental and biological changes. Some of the studies also shows that even a minimal exposure to EDC alters the normal functioning of the body leading to major health related issues⁸.

BPA as EDC

Action of BPA is at the hormonal level. When there is a distortion of hormonal balance, estrogenic related receptors are stimulated and brings about an abnormal estrogenic effect. Thus, giving rise to several hormone related abnormalities that has been previously reported in some studies (Fig:1)⁹. Similarly, BPA also acts on the endocrine system through several receptor mediated mechanisms. BPA acts like Xenoestrogen which binds and activates the estrogen receptor. Even though BPA has a lower affinity for genomic ER than estradiol, the circulating concentrations of BPA are biologically active and has higher affinity than estradiol. In addition, BPA is as bioactive as estradiol, particularly in those mediated by nongenomic signaling¹⁰.

Mechanism of BPA over some of the endocrine functions are as follows:

- BPA being an antiandrogen binds to androgen receptor and interferes the normal action of androgens.
- BPA alters the synthesis and circulation of steroid hormone concentrations

- Alters the thyroid activator receptor and
- Disrupts the glucocorticoid signaling ¹¹.

BPA acts on Estrogen Receptor (ER)

BPA can behave as a weak inducer of estrogen activity or also have an anti-estrogen effect. It can bind with ER α and ER β (estrogenic receptors). BPA blocks estrogenic response by competing with endogenous E2 ¹.

BPA as an Antiandrogen

Androgens act through a ligand-activated steroid hormone receptor called Androgen Receptor (AR). BPA acts by antagonising the AR signalling to inhibit the proliferation of Sertoli cells and male reproductive toxicology ¹⁴.

BPA alters steroid synthesis

BPA is an endocrine disruptor. Adult female exposure to BPA can cause potential effects on fertility, leads to a decrease in oocyte fertilization rate for women in vitro fertilization. BPA affects granulosa cells both in vivo and female fertility. Higher dose of BPA shows granulosa steroidogenesis, ovarian disruption and maximal reproductive potential. The effect of BPA on Follicle Stimulating Hormone stimulating the process of relatively high concentrations with human luteinized granulosa cells shows more sensitivity than KGN cell line ¹³.

BPA disrupts the Thyroid Hormone (TH) signalling

BPA acts on Thyroid Hormone in a negative feedback mechanism by inhibiting β -TR (Thyroid Receptor) action and involved in T4 elevation by the action of unopposed α -TR in the hippocampus. During Maternal exposure, BPA increases serum T4 and also shows a higher level of expression of RC3/Neurogranin. Eventual exposure of BPA may influence signalling of Thyroid Hormone (TH) in the developing fetal brain ¹⁵.

Developmental and parental exposure to BPA alters the cell morphology and embryological development of adult health and disease. It is associated with a decrease in young adult's anogenital distance which acts as an androgen-sensitive endpoint. BPA doesn't cause DNA mutation directly but induces epigenetic embryological changes ending in adult-onset disease ¹¹.

BPA as an Epigenetic Regulators

Epigenetic regulators play an important role in controlling gene expression by chromatin modification. Some of the epigenetic regulators are histone acetyltransferases, chromatin remodelling enzymes, methyltransferases etc., BPA induces epigenetic modifications in both human and animal cells. BPA not only modifies the methylation pattern of gene coding protein but also directly induces the gene responsible for DNA methylation (Fig: 2). BPA induces epigenetic modifications such as:

- a) DNA Methylation
- b) Histone Modification
- c) Non – Coding RNAs⁷

DNA Methylation

DNA methylation is a regular biological process of the epigenetic regulator. It is defined as methyl groups (CH₃) are added to the DNA molecule. DNA methylation process alters the DNA segment activity without changing the sequence (Fig:3). Transcription of Gene repression acts by DNA methylation¹⁶. DNA methylation either regulates the gene expression by recruiting the protein or inhibiting the transcription factors binding with DNA^{17,24}.

DNA methylation is essential for general development in human beings. It consists of following processes including

- Genomic Imprinting
- Inactivation of X-chromosomes
- Repression of transposable elements
- Ageing and
- Carcinogenesis

There are four bases in DNA. Only two bases of DNA can be methylated i.e., Cytosine and Adenine. DNA base with addition of methyl group to cytosine is commonly seen in all organisms⁶. In plants and other environmental organisms, DNA methylation is found in three different sequences:

- CG
- CHG or
- CHH

In mammals, DNA methylation is usually found in CG dinucleotides with the cytosines on both strands methylated²⁶. Non – methylated CG found in embryonic stem cells, neural tube development and hematopoietic progenitor cells¹⁷.

DNA methylation disrupts the gene transcription in two ways.

1. Binding of transcriptional proteins to gene are prevented or delayed
2. Methyl CpG – binding domain proteins (MBDs) – Protein bound methylated DNA¹⁷.

MBD protein includes additional proteins to the gene locus like histone deacetylases and other remodelling chromatin proteins that can modify histones and forming compact, inactive form of chromatin called as heterochromatin¹⁶.

DNA methylation is a heritable epigenetic alternator which causes carcinoma, atherosclerosis, genomic imprinting disorders and cardiovascular diseases¹⁸.

There are three major methods to identify and quantify DNA methylation.

1. Conversion and sequencing of Sodium bisulfite
2. Differential enzymatic DNA cleavage and
3. Affinity captures of methylated DNA.

Restriction enzyme based differential cleavage of methylated DNA is locus-specific.¹⁶

Histone modification

A histone is a protein that helps to form chromatin structure (DNA-wrapped protein octamers). Histone methylation is one of the post-translational epigenetic modification. It is defined as transfer of methyl groups to histone proteins with the help of histone methyltransferases (HMTs)¹⁹. Histone modifications include covalent modifications such as addition or removal of chemical groups such as acetyl groups, methyl groups, phosphates, and ubiquitin to the positively charged histone protein. These modifications change the chromatin structure by alteration of gene expression²⁰.

Addition of acetyl groups to any of the DNA base (Lysine) in histone neutralizes the cations and loosening the DNA nucleosome's grip. Histone modifications proceeds the transcriptional process to detect DNA and genes are active²¹.

Three types of histone modification involved in chromatin remodelling. It primarily occurs at N- terminal histone tails. These enzymatic modifications are

- Acetylation
- Methylation
- Phosphorylation and
- Ubiquitination.

Non-Coding RNA (ncRNA)

A non-coding RNA are group of RNAs, originally does not encode functional proteins and regulates the expression of genes at post transcriptional and transcriptional level.

The involvement of Non-Coding RNA in Epigenetic modifications can regulate gene expression and controlling the cell differentiation in chromosomal level. The ncRNAs which are involved in epigenetic process are classified into two major groups:

1. Small or Short non-coding RNA and
2. Long non-coding RNA²².

Small or Short ncRNA

Small or Short ncRNAs plays major role in eukaryotes. Different Short ncRNAs are essential for developmental process. Different types of small coding ncRNAs are:

- MicroRNAs (miRNAs)
- Short Interfering RNAs (siRNAs)
- Piwi-interacting RNAs (PiRNAs)^{23,25}.

MicroRNAs (miRNAs)

MicroRNAs (miRNA) is an endogenous single stranded RNAs highly prone to structural changes. It is the expression product of biological gene. MicroRNAs can regulate hundreds of different gene at same time. It binds with specific mRNA to induce cleavage and degradation²².

Short interfering RNAs (siRNAs)

Short interfering RNAs (siRNA) is exogenous long double stranded RNAs. Its origins from viral infection either by gene transfer or gene target. Short interfering RNAs have similar action mechanism like miRNAs by mediating gene silencing and degradation²².

Piwi-interacting RNAs (piRNAs).

Piwi-interacting RNAs (piRNA) are named due to their specific reflection facts with the piwi protein family. It comes from single chain precursor and arising from long single stranded molecule. PiRNAs serve as a primary transcript. It has the ability to induce euchromatin histone modifications. PiRNA/Piwi protein acts as a specific determinants of DNA methylation in germ cells and mediates the methylation of transposons²².

Long ncRNAs

lncRNAs combine with modified chromatin proteins and recruit their catalytic activity to specific binding sites in the genome by chromatin modifications and induces gene expression. The majority of non-coding RNA transcription belong to the group lncRNAs. Long ncRNAs plays an essential role in process of dosage compensation, genomic imprinting, developmental differentiation and patterning and stress response. It can either upregulate or downregulate gene expression in all human beings²².

Conclusion

Endocrine system plays an important role in balancing hormone in our body. Reproductive glands and thyroid gland secreting hormones function together to maintain homeostasis^[14]. Many recent studies proved that indicating Endocrine Disrupting Chemical (EDC) causes developmental disorders, metabolic syndrome, obesity, cancer and infertility. Bisphenol A acts as an EDC causing developmental anomalies due to its Endocrine disrupting properties. Bisphenol A interferes with tissue and organ development, alters the mechanism of different diseases. Many researchers are effectively working to understand more about Endocrine Disrupting Chemicals and its impact on human health and environment.

References:

1. Marta Diana Komarowska, Adam Hermanowicz, Urszula Czyzewska, Robert Milewski, Ewa Matuszcza, Wojciech Milytk and Wojciech Debek. Serum Bisphenol A Level in Boys with Endocrine Cryptorchidism: A Step to Male Infertility? *International Journal of Endocrinology*. 2015; 8.
2. Kangovou article. How does BPA affect the environment? 2019 January.<https://www.kangovou.com/the-importance-of-eco-friendly-products/>
3. Steven Siracusa, Lei Yin, Emily Measel, Shenuxan Liang, Xiaozhong Yu. Effect of Bisphenol A and its analogs on reproductive health: A mini review. *Reproductive Toxicology*. 2018 August; 79: 96-123.
4. Dinesh Tiwari and Geeta Vanage. Bisphenol A Induces Oxidative Stress in Bone Marrow Cells, Lymphocytes, and Reproductive Organs of Holtzman Rats. *International Journal of Toxicology*. 2017; 36(2): 142-152.
5. Stephen J. Genuis, Sanjay Beesoon, Detlef Birkholz and Rebecca A. Lobo. Human Excretion of Bisphenol A: Blood, Urine, and Sweat (BUS) Study. *Journal of Environmental and Public Health Volume*. 2012 January; 1-10.
6. Lisa Jones, Fiona Regan. *Encyclopedia of Analytical science*. Endocrine Disrupting Chemicals, *Microchemical Journal* 2013;3 Available from: science direct.com
7. Karina Caballero-Gallardo, Jesus Olivero-Verbela and Jennifer L. Freeman. Toxicogenomics to Evaluate Endocrine Disrupting Effects of Environmental Chemicals Using the Zebrafish Model. *Current Genomics*. 2016 April; 17(6): 515-527.
8. National Institute of Environmental Health Sciences. Bisphenol A (BPA). Available from: <https://www.niehs.nih.gov/health/topics/agents/sya-bpa/index.cfm>
9. Okugbe E. Ohore, Songhe Zhang. Endocrine disrupting effects of bisphenol A exposure and recent advances on its removal by water treatment systems. A review. *Research Gate. Scientific African* 5. 2019 August; 1-12.
10. Migliaccio.M. Chioccarelli,C. Ambrosino,A. Suglia,F. Manfrevola,O. Carnevali,S. Fasano, R. Pierantoni. Characterization of Follicular Atresia Responsive to BPA in Zebrafish by Morphometric Analysis of Follicular Stage Progression. *International Journal of Endocrinology*. 2018 December
11. Jakub Kwintkiewicz, Yoshihiro Nishi, Toshihiko Yanase and Linda C. Giudice. Peroxisome Proliferator–Activated Receptor- γ Mediates Bisphenol A Inhibition of FSH-Stimulated IGF-1, Aromatase, and Estradiol in Human Granulosa. *Environmental Health Perspectives*. 2010 March; 118 (3): 401-406
12. Hyun Ju Lee, Soma Chattopadhyay, Eun-Yeung Gong, Ryun Sup Ahn, and Keesook Lee. Antiandrogenic Effects of Bisphenol A and Nonylphenol on the Function of Androgen Receptor. *Toxicological Sciences*. 2003 May; 75: 40-46.
13. David Melzer, Lorna Harries, Riccardo Cipelli, William Henley, Cathryn Money, Parll Mc Cormock, Anita young, Jack Guralnik, Luigi ferrucci, Stefanie Bandinelli, Anna Maria Corsi, Tamara Galloway. *Environmental health Perspectives*, 2011 December; 119 (12).
14. Endocrine Disruptors. National Institute of Environmental Health Sciences. Available from: <https://www.niehs.nih.gov/health/topics/agents/endocrine/index.cfm>

15. Kenji Moriyama, Tetsuyu Tagami, Takashi Akamizu, Takeshi Usui, Misa Saijo, Naotetsu kanamoto, Yuji Hataya, Akira Shimatsu, Hideshi Kuzuya, kazuwa Nakao. Thyroid Hormone is disrupted by Bisphenol A as an antagonist. *Journal of Endocrinology Metabolism*.2002 November; 87 (11):5185-90.
16. Lisa D Moore, Thuc Le, Guoping Fan. DNA methylation & its basic function. *Neuropsychopharmacology*. 2013 January; 38 (1): 23-38.
17. Hyun Sik Jang, Woo Jung Shin, Jeong Eon Lee, Jeong Tae Do. CpG and Non – CpG Methylation in epigenetic gene regulation and brain function. *Genes (Basel)*. 2017 June; 8 (6):148
18. Terumichi Nakagawa, Masatoshi wakui, Tetsu Hayahida, Chiyoko Nishime, Mitsuru Murata. Intensive Optimization and evaluation of global DNA methylation quantification using LC-MS/MS. *Analytical and Bioanalytical Chemistry*. 2019 October; 411: 7221-7231.
19. Peter Cheung, Priscilla Lau. Epigenetic regulation by histone methylation and histone variants. *Molecular Endocrinology*. 2005 March; 19 (3) :563-573.
20. Chafik Ghayor, Franz E. Weber. Epigenetic regulation of Bone remodeling and its impacts in osteoporosis. *International of Molecular science*. 2016 September; 17 (9): 1446.
21. Bilal Alaskhar Alhamwe, Razi Khalaila, Johanna Wolf, Verena Von Bulow, Hani Harb, Fahd Alhamdan, Charles S. Hii, Susan L. Prescott, Antonio Ferrante, Harald Renz, Holger Harn, Daniel P. Potaczek. Histone modifications and their role in epigenetics of atopy and allergic diseases.*Allergy, Asthma and Clinical Immunology*. 2018 May; 14:39.
22. Frances Xin, Lauren M. Smith, Martha Susiarjo, Marisa S. Bartolomei and Karl J. Jepsen. Endocrine-disrupting chemicals, epigenetics, and skeletal system dysfunction: exploration of links using bisphenol A as a model system. *Environmental epigenetics*. 2018 February; 4 (2): 1-12.
23. Mallikarjuna Rao Gedda, Piyoosh Kumar Babele, Kulsoom Zahra, Prasoon Madhukar. "Epigenetic Aspects of Engineered Nanomaterials: Is the Collateral Damage Inevitable?", *Frontiers in Bioengineering and Biotechnology*, 2019 September; 7.
24. Zachary M. Laubach, Christoper D.Faulk, Andres Cardenas. Nutrition, DNA Methylation and developmental origins of cardiometabolic disease: A signal systems approach. *Handbook of Nutrition, Diet and Epigenetics*. 2019 January; 43-60.
25. Qudeer Ahmed Abdul, Byung Pal Yu, Hae Young Chung, Hyun Ah Jung, Jae Sue Choi. Epigenetic modifications of gene expression by lifestyle and environment. *Archives of Pharmacal Research*. 2017 October; 40: 1219- 1237.
26. Billian Jin, yajun Li, D. Robertson. DNAMethylation. *Genes Cancer*. 2011 June; 2 (6): 607-617.